

Applied Environmental Research



Journal homepage: http://www.tci-thaijo.org/index.php/aer

Natural Radioactivity in Groundwater in Phra Nakhon Si Ayutthaya Province

Paradee Kodcharin¹, Udorn Youngchuay², Sopa Chinwetkitvanich^{1,*}

 ¹Department of Sanitary Engineering, Faculty of Public Health, Mahidol University Bangkok 10400, Thailand
² Thailand Institute of Nuclear Technology (Public Organization), NakornNayok 26120, Thailand
* Corresponding author: Email: sopa.chi@mahidol.ac.th; Phone: +668-1173-5668

Article History

Submitted: 16 August 2017/ Accepted: 16 October 2017/ Published online: 1 August 2018

Abstract

This research aims to study the specific activity of natural radioactivity in groundwater samples taken in the area of Phra Nakhon Si Ayutthaya Province, Thailand. Totally, sixty groundwater samples collected from wells in eight districts were monitored and determined for radionuclides ²²⁶Ra, ²³²Th and ⁴⁰K and gross alpha and beta were determined using high resolution gamma spectrometer and Canberra Tennelec Series 5 gas flow proportional counter, respectively. Most of the sixty wells serve for water consumption and some other wells for agricultural. The results showed that the average activity concentrations of the gross alpha and beta were 0.01±0.007 and 0.15±0.02 Bq/L and the specific activity of ²²⁶Ra, ²³²Th and ⁴⁰K were averagely 0.77±0.13, 1.03±0.19 and 15.56±1.28 Bq/L, and respectively. The activity concentrations of the gross alpha and beta and the specific activity of the radionuclides in these samples exhibited quite low as compared to the recommended reference level for human consumption reported by World Health Organization (WHO).

Keywords: Natural radioactivity; Groundwater; Phra Nakhon Si Ayutthaya Province

Introduction

Water is the most important source for life and makes up 70-75% of total body weight. While 70% of the world's surface is covered by water, only 0.3% of the total water resources on earth is drinkable and suitable for daily use [1]. For instance, human uses of water are agriculture, industry, consumption, household uses and recreation. Natural disasters have been more severe than before due to environmental balance is gradually destroyed. Significant reduction of rainfall and longtime of dry spells directly affect agricultural and industrial sectors, as well as water consumption. Groundwater resource is an alternative providing leverage to tackle a shortage of water supply in many areas. Groundwater is acknowledged about its risk of natural contamination from mineral dissolution of their host rocks, as well as anthropogenic contamination caused by human activities. Moreover, natural radioactive materials are also concerned, especially using groundwater as water supply in communities. Consumption of contaminated groundwater may cause adverse health effects.

The radioactivity in groundwater occurs mainly from radionuclides of the natural decay chains of ²³⁸U, ²³²Th and ⁴⁰K. Some radionuclides can be easily dissolved in water depending on mineralogical and geochemical composition of soil and rock, redox condition and residence time of groundwater transporting through soil and bedrock, and reaction of groundwater with soil and bedrock [2]. When the decayed radionuclides are taken into human body through ingestion or inhalation, radionuclides will cause internal exposure [3]. Ingested radionuclides are absorbed into blood and accumulated in specific tissues causing damages [4]. Some radioactive materials can cause toxicity to kidney and increase risk of cancer.

Due to data of radioactive concentrations in groundwater in Thailand are quite rare, this study is to monitor natural radioactivity occurrence in groundwater located in Phra Nakhon Si Ayutthaya Province, Thailand. The objective of this study is to evaluate the gross alpha and beta activity concentrations and natural radionuclides activity concentration levels (226Ra, ²³²Th and ⁴⁰K) for consideration. The gross alpha and beta activity concentrations and radionuclides specific activity were determined using Canberra Tennelec Series 5 gas flow proportional counter and high resolution gamma spectrometer, respectively. In addition, the observed data in this study was useful for evaluation of radioactive contamination level in these groundwater samples.

Material and methods

1) Monitoring design

The study area in this work is located in Phra Nakhon Si Ayutthaya province, 76 km

northern distance from Bangkok, covering an area of 2,556 km² and dividing into sixteen districts. This province is located in the flat river plain of the Chao Phraya river valley. The presence of four rivers, i.e. Chao Phraya River, Pa Sak River, Lopburi River and Noi River, flowing through the city makes this province a major rice farming area of Thailand. Due to drought phenomena have recently occurred, groundwater is widely used throughout the province as an alternative water reserve. Groundwater in this province serves for both consumption and agricultural purposes. Sixty wells were designnated for groundwater samples collection during May to October, 2016 as illustrated in Figure 1. Most of designated wells serve for consumption purpose through community water supply system. At least two liters of groundwater were taken for each sampling. Then, they were preserved with 1N HCl to pH 2 in order to avoid loss of radionuclide fraction by ad-sorption with the container and to prevent some biological activities according to EPA methods of 900.0 and 901.1.

2) Determination of gross alpha and beta radioactivity

One liter each sample was filtered via filter paper (Whatman No.1) and put into two-liter beaker. The filtrate with certain volume was further evaporated on hot plate until almost dry and its residue appeared. Then, residue was transferred to clean planchet and further drying to evaporation under infrared lamp. After drying process, residue was weighted and stored in desiccator until analytical measurement. Weighted residues were analyzed for the radiation (in terms of gross alpha and beta) by Canberra Tennelec Series 5 gas flow proportional counter with counting time of 120 minutes per sample. In addition, the control sample (or blank sample) was required and had to be placed in the front sequence of sample series during the measurement. The gross alpha and beta counting system were calibrated

using planchets containing of 241 Am and 90 Sr/ 90 Y standard sources, respectively. The counting efficiencies for the system were 23% for alpha and 36% for beta. Finally, the measurement result was used to calculate for radioactivity with the following equations (Eq. 1) as shown below.

3) Determination of radioactivity of radionuclides (²²⁶Ra, ²³²Th and ⁴⁰K)

One liter of each water sample was transferred into a Marinelli beaker. The beakers was subsequently firmly sealed for at least four weeks to ensure a state of secular equilibrium between radium isotopes and their respective daughters radionuclides measuring of gamma radiation [5]. The sample was analyzed for ²²⁶Ra, ²³²Th and ⁴⁰K using a high resolution gamma spectrometer. The efficiency calibration of the detector was performed using multinuclide distributed in 1.0g/cc epoxy matrix calibration standards with approximate volume of 1 liter in a Marinelli beaker. The efficiency calibration of the detector was performed by Cs-137 and Co-60, which the energy peak of Cs-137 was 662 keV and those of Co-60 were 1173 keV and 1333 keV. Each water sample in a Marinelli beaker was determined for radionuclides using high resolution gamma spectrometer counting for 50,000 seconds. A result of which was further calculated for radioactivity with the equations (Eq. 2) as shown below.

4) Annual dose calculation

In this study, annual dose ingestion of 226 Ra contamination in groundwater was considered. Therefore, the radioactivity dose from annual consumption in each adult was estimated according to the following equation (Eq. 3) [6]; considering with consumption rate of 2 liters/ day and the conversion factors of 0.28 µSv/Bq.



Figure 1 Sampling sites in Phra Nakhon Si Ayutthaya province.

Activity concentration (Bq/L) = $\frac{(A_s - A_b) \times 100}{\% \epsilon \times V \times 60}$ (Eq. 1)

Where: A_s is count rate of samples (cpm), A_b is count rate of background (cpm), $\Re \epsilon$ is efficiency percentage of the detector, V is volume of water (L) and 60 is conversion factor from dpm/Bq.

Activity Concentration (Bq/L) =
$$\frac{N_{(cps)}}{\epsilon x V x P_{v}(E)}$$
(Eq. 2)

Where: $N_{(cps)}$ is net counts of radionuclide in the sample (count per second), ε is the efficiency of energy detector, V is volume of water (liter) and $P_{\gamma}(E)$ is opportunity to decay and emit gamma ray energy.

Effective dose
$$(mSv/y) = A_w x IR_w x ID_F$$
 (Eq. 3)

Where: A_w is activity concentration (Bq/L), IR_w is intake of water for person in 1 year (730 L/year) and ID_F is the effective dose equivalent conversion factor (μ Sv/Bq)

Results and discussion

ł

The activity concentrations of gross alpha and beta and specific activity of the radionuclides (226 Ra, 232 Th and 40 K) in groundwater samples in Phra Nakhon Si Ayutthaya Province were concluded in Table 1. From the results, pH values of sixty samples were observed in the range of 6.7-8.4, most of which were in the neutral pH with the average of 7.6. Their conductivities were in the range of 232 - 3946 µc/cm where their average value was 902 µc/cm).

The activity concentrations of gross alpha were found in the range of 0.009-0.029 Bq/L, as well as those of gross beta were in the range of 0.063-0.345 Bq/L. The averages of activity concentration of gross alpha and beta observed in this area were 0.011±0.007 and 0.156±0.018 Bq/L, respectively, which were comparatively lower than their recommended reference levels by WHO of 0.5 and 1 Bq/L, respectively (Figure 2). Especially, the maximum concentration of gross alpha (0.029 Bq/L) was considered as only 6% of the reference level (0.5 Bq/L). However, the maximum concentrations of gross beta (0.345 Bq/L) were in considerable percentage (about 35%) of the reference level of 1 Bq/L.

1) Radium-226

The concentrations of radionuclides ²²⁶Ra, ²³²Th and ⁴⁰K were found in the ranges of 0.520-0.921, ND-1.033 and 12.9-18.8 Bq/L, respectively (Figure 3). Their averages of activity concentrations were 0.77 ± 0.13 , 1.03 ± 0.19 and 15.6 ± 1.29 Bq/L, respectively. In case of ²²⁶Ra, their activity concentrations were observed in the range that its maximum concentration (0.921 Bq/L) was much closed to the recommended level of 1 Bq/L for water consumption announced by WHO.

Table 2 gathered several reports of radioactivity monitoring in various areas in order to compare with our study. Similarly, Zhuo et al. (2001) [7] reported the maximum value of 0.93 Bq/L was found in groundwater sample taken in Fujian province (Table 2). Nevertheless, some samples contained higher concentrations of ²²⁶Ra were observed in groundwater samples taken in Egypt [8] and Yemen [9] and Serbia [10].

groundwater samples collected in Phra Nakhon Si Ayutthaya Province during May to October, 2016										
Parameter		Concentration			Standard deviation					
	Max	Min	Average	Max	Min	Average				
Gross Alpha (Bq/L)	0.029	0.009	0.011	0.010	0.004	0.007				
Gross Beta (Bq/L)	0.345	0.063	0.156	0.024	0.013	0.018				
²²⁶ Ra (Bq/L)	0.921	0.520	0.767	0.166	0.110	0.133				
²³² Th (Bq/L)	1.033	ND	-	-	-	0.194				
⁴⁰ K (Bq/L)	18.8	12.9	15.6	3.635	0.912	1.290				
pH	8.4	6.7	7.6	-	-	-				
Conductivity (µc/cm)	3946	232	902	-	-	-				

Table 1 Activity concentration (Bq/L) of natural radioactivity and associated characteristics in groundwater samples collected in Phra Nakhon Si Ayutthaya Province during May to October, 2016

ND is Not Detectable



Figure 2 Interpolated radiological maps for gross alpha and beta in Phra Nakhon Si Ayutthaya Province.



Figure 3 Interpolated radiological maps for ²²⁶Ra and ⁴⁰K in Phra Nakhon Si Ayutthaya Province.

Country	Type of	Activity concentration (Bq/L)					Ref.
	water	Alpha	Beta	²²⁶ Ra	²³² Th	⁴⁰ K	
Turkey	Groundwater	0.08-0.38	0.12-3.47	-	-	-	[2]
Hungary	spring waters	0.03-1.75	0.03-2.01	-	-	-	[11]
Bangladesh	Surface water	0.45x10 ⁻³ - 1.36x10 ⁻³	0.06-0.28	-	-	-	[12]
Nigeria	Underground water	0.31-14.49	0.02-27.5	-	-	-	[13]
China	Groundwater	-	-	0.001-0.94	-	-	[7]
Roi-Et Province	Groundwater	-	-	<0.006-0.177	-	-	[14]
Egypt (Elba)	Groundwater	-	-	1.6-11.1	0.21-0.97	9.1-23	[8]
Yemen	Groundwater	-	-	0.86-3.09	0.46-2.01	7.84-18.02	[9]
Jordan	Hot spring water	-	-	3.8–6.8	1.42–2.37	23.2-34.8	[15]
Serbia	Groundwater	0.001-1.33	0.02-5.43	0.005-2.56	0.006-0.79	0.012-2.6	[10]
Phra Nakhon Si Ayutthaya	Groundwater	0.009- 0.029	0.063- 0.345	0.520- 0.921	N.D1.033	12.9-18.8	Present work

Table 2 The activity concentration of natural radioactivity in Bq/L of water samples reported in various studies.

However, there is a report of ²²⁶Ra monitoring in groundwater samples taken in Roi-Et province, northeastern part of Thailand [14]. Those samples found in the range of <0.006-0.177 Bq/L, which were lower than in our study. This was probably due to different geology constituting water-bearing formations of groundwater system. Geological map of Thailand 1: 250,000 of Phra Nakhon Si Ayutthaya province [16] illustrated some of igneous rocks in its geology, but there was none shown in that of Roi-Et province [17]. Some literature reviews mentioned that ²²⁶Ra and ²²⁸Ra levels in groundwater related to certain types of rock (e.g. granite, sandstone) or rocks dissolution [18]. Anyway, some admitted that wide ranges of radioactivity were observed with little correlation to the type of rock or sediment constituting aquifer formation [19].

2) Thoriun-232

In case of ²³²Th (radiological maps were not shown here), there was no data of ²³²Th concentrations in the study of groundwater in Roi-Et province [14]. Usually, thorium content in crustal rocks is observed at low level. In addition, thorium typically forms complex compounds with ions in water either acidic or basic conditions. ²³²Th concentrations observed in this study were averagely 1.0 ± 0.19 Bq/L, similar range of which was reported in groundwater samples collected in Yemen [9] and hot spring water samples in Jordan [15]. In contrast, the much lower concentrations of ²³²Th in groundwater were mentioned in samples taken in Serbia [10].

3) Potassium-40

The ⁴⁰K concentrations monitored in this study were in moderate level in comparison to elsewhere [8-10, 15]. Theoretically, the abundance of ⁴⁰K activity observed in groundwater was usually explained by the relevance to potassium fertilizer application in agricultural activities and transportation through groundwater system. Anyway, higher concentrations of ⁴⁰K were found in hot spring water samples studied in Jordan [15]. Similar concentrations were observed in groundwater samples monitored in Yemen [9]. Much lower concentrations were reported in a study of groundwater in Serbia [10]. It is possible that moderate concentrations of ⁴⁰K activity found in this study area might be resulted from potassium fertilizer application. More statistical data of background level of ⁴⁰K activity and fertilizer application in Phra Nakhon Si Ayutthaya province are required for in depth discussion.

4) Annual dose of ²²⁶Ra ingestion

The annual dose calculation for adults, considering only ingestion from ²²⁶Ra by the water consumption rate is 2 L/day in this study were found in the range of 0.106-0.188 mSv/ year. Although ²²⁶Ra level found in groundwater samples taken in Phra Nakhon Si Ayutthaya Province during May to October, 2016 (Table 2) complied with WHO guideline of 1 Bq/L, all results of annual dose calculation exceeded the individual dose criteria (IDC) of 0.1 mSv/yr suggested by WHO (2011) [20]. However, this screening level for drinking water usually applied if either gross alpha activity or gross beta respectively exceeded 0.5 and 1 Bq/L guidance concentrations, which those in this study were not in the case (Table 2). Anyway, this study attempted the annual dose calculation for ²²⁶Ra in order to compare with the study in Roi-Et province [14]. Thus, the annual doses of ²²⁶Ra in our study were higher than those in Roi-Et province due to higher ²²⁶Ra concentration as previously mentioned. The most frequency of 38% (of total 60 data) were observed in range of 0.101-0.120 mSv/yr as shown in Figure 4. Nevertheless, WHO [20] also suggested that this IDC (0.1 mSv/yr) should be determined only in case of any radionuclide was to exceed the guideline concentration; consequently resulting in sumthe continuation monitoring for a whole year water is unsuitable for consumption.

mary annual dose might exceed the IDC. Then, should be applied before concluding that the



Figure 4 Frequency distribution of effective dose for ²²⁶Ra in groundwater in water sample.

Conclusion

The natural radioactivity levels of gross alpha and beta and radionuclides: ²²⁶Ra, ²³²Th and ⁴⁰K observed in groundwater samples taken in Phra Nakhon Si Ayutthaya Province during May to October 2016 considerably complied with the guidance level for drinking water quality recommended by WHO. The ²²⁶Ra levels in this area were in concern where their annual dose calculation resulted in exceeding the individual dose criteria adopted by WHO (2011). However, this was too early to conclude these groundwater samples were unsuitable for drinking purpose. Different seasonal monitoring will be further studied and their results will be compared and discussed with data presented here. Therefore, the annual dose for ingestion of the radionuclides found in groundwater will be more evaluated and discussed.

Acknowledgements

The authors express their thanks to the Thailand Institute of Nuclear Technology (Public Organization) for laboratory assistance and support. The authors would like to thanks the Department of Sanitary Engineering, Faculty of Public Health, Mahidol University for all supports.

References

- [1] Althoyaib, S.S., El-Taher, A. Natural radioactivity measurements in groundwater from Al-Jawa, Saudi Arabia. Journal of Radioanalytical and Nuclear Chemistry, 2015, 304, 547-552.
- [2] Turhan, Ş., Özçitak, E., Taşkin, H., Varinlioğlu, A. Determination of natural radioactivity by gross alpha and beta measurements in ground water samples. Water Research, 2013, 47, 3103-3108.
- [3] Kobya, Y., Taskin, H., Yesilkanat, C.M., Çevik, U., Karahan, G., Çakır, B. Radioactivity survey and risk assessment study for drinking water in the ArtvinProvince, Turkey. Water Air and Soil Pollution 2015, 226(3), 1-9.
- [4] Canu, I.G., Laurent, O., Pires, N., Laurier, D., Dublineau, I. Health effect of naturally radioactivity water ingestion: The need for enhanced studies. Environmental Health Perspectives, 2011, 119, 1676-1680.
- Ibrahim, M., Shalabiea, O., Diab, H. Mea-[5] surement of some radioactive elements in drinking water in Arar city, Saudi Arabia. American Journal of Life Sciences, 2014, 2(1), 24-28.

- [6] Harb, S., El-Kamel, A.H., Zahran, A.M., Abbady, A., Ahmed, F.A. Natural radioactivity of ground water in some areas in Aden Governorate South of Yemen Region. Proceedings of the 1st International Conference on New Horizons in Basic and AppliedScience, Hurghada - Egypt, 2013, 1(1), 281-289.
- [7] Zhuo, W., Iida, T., Yang, X. Occurrence of ²²²Rn, ²²⁶Ra, ²²⁸Ra and U in groundwater in Fujian Province, China. Journal of Environmental Radioactivity, 2001, 53, 111-120.
- [8] El Arabi, A.M., Ahmed, N.K., Salahel, D.K. Natural radionuclides and dose estimation in natural water resources from Elba Protective area, Egypt. Radiation Protection Dosimetry, 2006, 121(3), 283-292.
- [9] Saleh, E.E., El-Mageed, A.I.A., El-Gamal, H., Hussien, M.T. Assessment of radiation hazards a result of natural radio-activity in water fromAbyan delta, Yemen. Journal of Radioanalytical and Nuclear Chemistry, 2015, 304, 1235-1241.
- [10] Ćuk, M., Papić, P., Stojković, J. Natural radioactivity of groundwater in Serbia. AnnalesGéologiques De La Péninsule Balkanique, 2013, 74, 63-70.
- [11] Jobbágy, V., Kávási, N., Somlai, J., Dombovári, P., Gyöngyösi, C., Kovács, T. Gross alpha and beta activity concentration in spring waters in Balaton Upland, Hungary. Radiation Measurements, 2011, 46, 159-163.
- [12] Biswas, S., Ferdous, J., Begum, A., Ferdous, N. Study of gross alpha and gross beta radioactivities in environmental samples. Journal of Scientific Research, 2015, 7(1-2), 35-44.

- [13] Atsor, A.J., Akpa, T.C., Akombor, A.A. Determination of gross alpha and beta radioactivity in underground water at Gboko and its environs. Research Journal of Physical Sciences, 2015, 3(6), 1-9.
- [14] Quinram, P., Yenchai, C. Specific activity of radium-226 in groundwater in the area of Roi-Et Province. (In Thai). Thaksin University Journal, 2015, 18(3), 209-214.
- [15] Saqan, S.A., Kullab, M.K., Ismail, A.M. Radionuclides in hot mineral spring waters in Jordan. Journal of Environmental Radioactivity, 2001, 52, 99-107.
- [16] Geological Survey Division, Department of Mineral Resources. Geological Map of Thailand 1:250,000 Changwat Phra Nakhon Si Ayutthaya (sheet ND 47-8) 1985a. Printed by Royal Thai Survey Department.
- [17] Geological Survey Division, Department of Mineral Resources. Geological Map of Thailand 1:250,000 Changwat Roi Et (sheet NE 48-14) 1985b. Printed by Royal Thai Survey Department.
- [18] Godoy, J.M., Godoy, M.L. Natural radioactivity in Brazilian groundwater. Journal of Environmental Radioactivity, 2006, 85, 71-83.
- [19] Chau, N.D., Dulinski, M., Jodlowski, P., Nowak, J., Rozanski, K., Sleziak, M., Wachniew, P. Natural radioactivity in groundwater - a review. Isotopes in Environmental and Health Studies, 2011, 47 (4), 415-437.
- [20] World Health Organization. Guidelines for Drinking-Water Quality (fourth edition), 2011. [Online] Available from: http:// www.who.int/water_sanitation_health/ publications/2011/dwq_guidelines/en/ [Accessed 15 April 2017].