

# GENETIC VARIATIONS IN *AEROMONAS HYDROPHILA* ISOLATES FROM CLINICAL AND ENVIRONMENTAL SOURCES IN THAILAND

S Korbsrisate<sup>1</sup>, S Trakulsomboon<sup>2</sup>, S Damnin<sup>1</sup>, J Gatedee<sup>3</sup>, OA Ratchtrachenchai<sup>4</sup>,  
A Leelaporn<sup>5</sup>, B Eampokalap<sup>6</sup>, C Gongviseisoog<sup>6</sup> and S Suparak<sup>1</sup>

<sup>1</sup>Department of Immunology, <sup>2</sup>Department of Medicine, <sup>5</sup>Department of Microbiology, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok; <sup>3</sup>Faculty of Medical Technology, Rangsit University, Pathum Thani; <sup>4</sup>Department of Medical Sciences, National Institute of Health, Nonthaburi; <sup>6</sup>Microbiology Laboratory, Bamrasnaradura Hospital, Nonthaburi, Thailand

**Abstract.** *Aeromonas hydrophila*, a widely distributed human pathogen causing a variety of diseases, can be isolated from clinical and environmental sources. Analysis in Thailand of 110 isolates of *Aeromonas hydrophila* by randomly amplified polymorphic DNA-PCR (RAPD-PCR) revealed one specific RAPD pattern group (G) that was associated only with strains from environmental sources. Cytotoxic activity, adhesion to epithelial cells and exoenzyme secretions of *A. hydrophila* were also investigated. A comparison of isolates with pattern group G with a set of isolates derived from human blood showed low induction of cytotoxicity from those with RAPD pattern group G suggesting low virulence of these strains.

---

Correspondence: S Korbsrisate, Department of Immunology, Faculty of Medicine Siriraj Hospital, Mahidol University, 2 Pran Nok Road, Bangkok 10700, Thailand.  
E-mail: grsks@mahidol.ac.th