

Research Article

Improvement of the quality and shelf-life of minced beef mixed with soy protein by Sage (*Saliva officinalis*)

Ali M. Ahmed^{1*} and Takwa H. Ismail²

¹Department of Food Hygiene and Control, Faculty of Veterinary Medicine, Suez Canal University, Ismailia, Egypt

²Department of Food Hygiene, Animal Health Research Institute, Ismailia Lab., Egypt.

*Email: meawad@maktoob.com

Abstract

The effect of various concentrations of sage (*Saliva officinalis*) on the microbial quality and shelf-life of minced beef mixed with soybean protein was investigated. For this purpose, minced beef mixed with soybean protein samples were divided into four groups. First group was kept as a control group, while other ones were treated with 0.1%, 0.3% and 0.5% sage extract. All samples were stored at 4°C and microbiological evaluation was conducted at intervals of 0, 1, 3, 5 and 7 days post-storage for aerobic, anaerobic, enterobacteriaceae, yeast-mould, *Bacillus cereus* and *Staphylococcus aureus* counts. Minced beef pose a risk of food-borne infection or intoxication due to existence and growth of *B. cereus* and *S. aureus*. Sage extract concentration of 0.3% and 0.5%, significantly reduced aerobic, anaerobic, enterobacteriaceae, yeast-mould, *Bacillus cereus* and *Staphylococcus aureus* counts in the samples. Control and 0.1% sage groups did not comply with the Egyptian Standards Criteria for total aerobic counts when compared to other groups within the seven days of shelf-life. The results indicated that 0.3% and 0.5% sage extract improved the microbiological quality and prolonged the shelf-life of the minced beef to seven days of retail displayed at 4°C.

Keywords: food additives, *Bacillus cereus*, *Staphylococcus aureus*, meat, spoilage, Egypt.

Introduction

Minced beef mixed with soybean is one of the meat products widely consumed in Egypt. Microbial quality of meat products plays an important role in increasing public health issues all over the world. During the last decade, there were great improvements in hygienic techniques for production of meat products with attention of a lot of consumers towards healthy nutrition. The use of antimicrobial ingredients is one of the widely used methods to maintain microbiological safety and prolong the shelf-life of food products [1].

According to the Egyptian Organization for Standardization and Quality Control [2], a minced beef mixed with soybean protein is defined as a fresh meat mixed with soybean protein with or without the addition of food salt or spices and stored at chilling or freezing condition. The use of soybean protein (food grade) as meat products extenders is widely spread in meat products industry because

of their interesting nutritional and functional properties. Together with these, health diets and economical reasons are the major causes for the addition of soybean proteins to meat products [3].

Herbs are used as substances enhancing the taste and varieties of regular food. Some herbs are used as meat additives reporting to have bactericidal or bacteriostatic activities [4]. The inhibitory effects of herbs are mostly because of the volatile oils [5, 6]. Sage, *Salvia officinalis*, is a perennial herb, with gray green aromatic leaves and their name is derived from the Latin word *salvere*, meaning "to save". Ancient Greek-Romans and ancient Egyptians used sage as meat preservative and as a medicinal herb. In the Middle Ages, it was used in treating cholera, common cold, fevers and epilepsy [7]. The variety used nowadays in cooking is flowerless and with broad leaves. Herbs have a very powerful flavour and can be used fresh or dried.

Minced beef is not only highly susceptible to spoilage, but also it is frequently involved in the spread of food-borne illness. During slaughtering and processing of cattle, all potentially edible tissues are subjected to microbial contamination from various sources. Aerobic plate counts often are chosen as an indicator of the effectiveness of hazard analysis critical control point (HACCP) plans, because data for all aerobic bacteria are more easily collected than data for pathogens of concern or other indicator organisms [8]. Enterobacteriaceae are very useful as indicators of bad hygiene or bad treatment of food products and their presence in large number indicates a good possibility of their multiplication, implying multiplication of other pathogens [9]. It is suggested that yeast and mould play an important role in causing spoilage of meat [10, 11]. *Bacillus cereus* and *Staphylococcus aureus* are two important food-borne pathogens. The most staphylococcal food poisoning and *B. cereus* food poisoning outbreaks may be due to poorly cooked and extensively handled meat products [12].

Several studies have revealed the antimicrobial properties of sage [13, 14, 15, 16, 17, 18], however, researchers did not lend their attention to the antimicrobial properties of sage for improving the microbial quality of minced beef mixed with soybean protein. Therefore, the objective of this study was to evaluate the efficacy of sage extract for improving the microbial quality and prolonging the shelf-life of minced beef mixed with soybean protein when stored at 4°C.

Materials and Methods

Preparation of sage extracts

The dried herbs, *Salvia officinalis*, obtained from local market at Ismailia city, Egypt were individually ground by an electric grinder to pass through 60 mesh sieve. About 5 g of ground herb was shaken for 30 min. with 50 mL of 96% ethanol at room temperature (22°C ±2). The mixture was allowed to stand for 24 hours. Then it was filtered to obtain alcoholic particle free sage extract.

Experimental design

Firstly, fresh beef was purchased from a local market at Ismailia city, Egypt on the day of preparation. The meat was then cut and minced with a grinder through a 4 mm plate diameter. Minced beef was mixed with 30% of their weight by soybean protein (Cargill, Food Grade) previously soaked in potable water for at least 4 hours. The samples were divided into four groups; each group included 25 samples, each one weighing 200 g. The four groups were thoroughly mixed in sterile mixer with 0% (control group), 0.1%, 0.3% and 0.5% sage extracts respectively. Each sample was wrapped with saran wrap and placed in a chiller at 4°C ±1. During chilling at intervals of 0, 1, 3, 5 and 7 days, five samples were removed for microbiological evaluation.

Sample preparation

Minced beef samples of 25 g were removed aseptically and homogenized in 225 mL of 0.1% sterile peptone water for 2 min using a Stomacher 400 Lab Blender to provide dilution of 10⁻¹. From the

original homogenate, 1 mL was transferred aseptically to a test tube containing 9 mL sterile 1% peptone water (w/v) to prepare dilution of 10^{-2} , then from which further ten fold decimal serial dilution up to 10^{-8} were prepared. From these serial dilutions, the bacteriological investigations were performed.

Microbiological evaluation

Appropriate diluents of each tube were placed on the following media in duplicate; Plate Count Agar (Difco Co., Ltd.) for total aerobic count; Reinforced Clostridial Medium (Oxoid, CM 0149) for total anaerobic count; Violet Red Bile Glucose Agar (Oxoid, CM 485) for total enterobacteriaceae counts; Malt Extract Agar (Oxoid, CM 0059) for total yeast and mould count; Mannitol Egg Yolk Polymixin Agar (Oxoid, CM 0929) for total *B. cereus* count and Baird-Parker Agar (Oxoid, CM 275) with Egg Yolk-Tellurite Emulsion (Oxoid, SR 54) for total *S. aureus* count.

Statistical analysis

Experimental trials were repeated twice. The statistical analysis for this study was performed using SAS for Windows. Data was analyzed for the significant differences due to sample treatment with different levels of sage extract and storage for different time periods. Data were subjected to one-way analysis of variance (ANOVA). Any significant differences ($P < 0.05$) were analyzed by the multiple comparisons procedure of LSD (least significant difference), using a level of significance of $\alpha = 0.05$.

Results and Discussion

Results are presented in Table 1. Addition of sage extract decreased the initial microbial count partially and slowed down the growth during the storage period parallel to the increasing concentration. In general, the microbiological quality of meat products as purchased by the consumer relies on a number of factors, such as the quality of the raw materials, other materials used or added during processing operations to the products as extraneous contaminants, sanitation during processing and packaging. At concentration of 0.3% and 0.5%, sage significantly reduced the mean aerobic counts in the samples. The bactericidal effects of 0.5% sage extract in laboratory media was determined on various microorganisms which are commonly encountered in the food industry [13, 18, 19].

EOS [2], set a standard maximum limit of 10^6 cfu/g for total aerobic count in minced beef mixed with soybean protein with maximum 7 days of storage at 4°C. Based on this standard, control groups and 2nd group (0.1% sage) did not comply with the Egyptian Standards criteria for that product. The most important effect of 0.3% and 0.5% sage extract on treated minced beef is a significant improvement in its shelf-life up to 7 days of storage at 4°C.

As sage extract concentration increased the total anaerobic bacteria counts decreased. This reduction was even faster in samples containing 0.3% and 0.5% sage extract; no growth of anaerobic bacteria was observed in the analysis after 1 and 3 days of storage for samples treated with 0.5% sage extract. Generally, the significantly lower mean anaerobic counts obtained for treated samples reflect the inhibitory effect of the sage extract, in addition to the effect of chilling temperature on the growth of anaerobic bacteria.

Enterobacteriaceae family includes facultative anaerobic gram negative straight bacilli, motile employ peritrichous flagella. Most members of this family are mainly mesophilic while some strains can grow at 0°C [20]. Enterobacteriaceae family include some food-borne genera such as *Escherichia*, *Salmonella*, *Klebsiella*, *Serratia*, *Enterobacter*, *Citrobacter*, *Yersinia*, *Proteus*, *Providencia*, *Shigella* and *Erwinia* that pose a great health risks for consumers [21].

Table 1. Effect of Different Concentrations of Sage Extract on Microbial Quality (Counts in log₁₀ cfu/g) of Minced Beef Stored at 4°C.

		Sampling Time (days)				
Sage extract, %	0	1	3	5	7	
Aerobes						
0	4.29 ^{a*}	4.77 ^a	5.64 ^a	6.74 ^a	6.78 ^a	
0.1	2.40 ^a	2.44 ^b	3.90 ^b	5.11 ^b	6.12 ^b	
0.3	1.96 ^b	1.40 ^{bc}	2.20 ^c	3.20 ^c	3.87 ^c	
0.5	1.64 ^c	0.55 ^d	1.57 ^d	2.04 ^d	3.58 ^{cd}	
Anaerobes						
0	1.39 ^{a*}	1.71 ^a	1.98 ^a	2.19 ^a	3.15 ^a	
0.1	1.18 ^{ab}	1.22 ^b	1.90 ^{ab}	2.10 ^{ab}	2.79 ^b	
0.3	0.80 ^c	0.97 ^c	1.18 ^c	1.27 ^c	1.50 ^c	
0.5	0.34 ^d	ND	ND	0.35 ^c	0.98 ^d	
Enterobacteriaceae						
0	2.37 ^{a*}	2.51 ^a	2.92 ^a	3.99 ^a	4.95 ^a	
0.1	1.11 ^b	1.12 ^b	1.90 ^b	2.81 ^b	3.35 ^b	
0.3	0.60 ^c	0.67 ^c	1.15 ^c	1.96 ^c	2.50 ^c	
0.5	0.32 ^d	0.20 ^d	0.87 ^d	1.85 ^{cd}	2.56 ^{cd}	
Yeast & Mould						
0	3.01 ^{a*}	3.51 ^a	3.52 ^a	3.60 ^a	3.67 ^a	
0.1	2.52 ^b	2.47 ^b	2.20 ^b	2.74 ^b	3.02 ^b	
0.3	1.84 ^c	1.20 ^c	2.11 ^{bc}	2.50 ^{bc}	2.59 ^c	
0.5	1.42 ^d	0.50 ^d	0.99 ^d	1.11 ^d	1.20 ^d	
B. cereus						
0	2.24 ^{a*}	2.67 ^a	2.84 ^a	3.66 ^a	4.68 ^a	
0.1	1.85 ^b	ND	0.76 ^b	1.90 ^b	2.78 ^b	
0.3	1.00 ^c	ND	ND	ND	ND	
0.5	0.57 ^d	ND	ND	ND	ND	
S. aureus						
0	0.84 ^a	1.21 ^a	1.84 ^a	2.14 ^a	2.63 ^a	
0.1	0.57 ^b	0.32 ^b	1.24 ^b	1.75 ^b	2.41 ^b	
0.3	0.21 ^c	0.13 ^c	ND	ND	ND	
0.5	0.20 ^d	ND	ND	ND	ND	

Means in the same column (a, b, c, d) and in the same row (p, q, r, s) with different letters for each microbial group differ significantly ($P < 0.05$).

ND = not detected

The results confirmed the significant ($P < 0.05$) inhibitory effect of different concentrations of sage extract combined with chilling temperature on the growth of enterobacteriaceae compared to control sample. 0.3% and 0.5% sage extract were more effective in lowering the enterobacteriaceae counts compared to 0.1% sage extract concentration.

Yeast and mould contribute a definite part of the natural microflora of meat. Their counts are used as an index of storability and sanitary quality of the products. The total yeast and mould counts of all sage treated groups were significantly lower than that of the control samples. Fungicidal activity of sage has been reported by Farag, *et al.*, [22] and Schmitz *et al.*, [23].

The presence of yeast and mould in meat products are objectionable, as they grow at a wide range of temperature and pH values, resulting in spoilage of the product [24]. Such yeast and mould might cause gas and off flavour in chilled food on account of their proteolytic activity, rancidity of cold-

stored meat products spoiled by yeast is mainly the cause of lipolytic activity by yeast [25]. Some pathogenic yeast in meat products make these products unsuitable for human consumption [26].

The addition of 0.3% and 0.5% of sage extract to minced beef was effective in inhibiting the growth of *B. cereus*. During slaughtering process, meat is exposed to various sources of *B. cereus* contamination.

Most gram positive bacteria were susceptible to the sage extract, but showed considerable variation in their extent of inhibition. Most of *Bacillus* species were killed at the 0.5% sage concentration, however, the minimum inhibitory concentrations of sage extract were 1.25-2.5 $\mu\text{L}/\text{mL}$ for *S. aureus* and 0.15-2.5 $\mu\text{L}/\text{mL}$ for *B. subtilis*. [14, 19, 27]. Sage at concentration 0.2% did not inhibit the growth of the majority of bacteria [28].

Staphylococcus species are Gram positive nonmotile, nonspore forming, facultative anaerobic cocci occurring in pairs or irregular clusters. The initial counts of *S. aureus* for control group were higher than treated groups. The inhibitory effect of sage extract against *S. aureus* has also been reported [14, 27].

The most important food-borne genera of staphylococci are *S. aureus* coagulase positive. The growth of *S. aureus* in food is a potential public safety hazard since many of its strains produce enterotoxins that cause food poisoning when ingested. *S. aureus* can be isolated from meat grinders, knives, saw blades and cutting boards or tables [20].

Conclusion

Microbiological control in minced beef has been identified as an important factor in improving quality, extending the shelf-life of the product and protecting consumers from the hazards of food-borne illness. The results indicate that the minced beef mixed with soybean when stored at 4°C are still contaminated with high levels of microorganisms. Sage extract was found to be significantly effective in improving microbiological quality and prolonging the shelf-life of the minced beef to day 7 of retail display.

References

1. Bİngöl, E. and Bostan, K. (2007): Effect of Sodium Lactate on the Microbiological Quality and Shelf Life of Sausages. *Turkish Journal of Veterinary Animal Science*, 31(5): 333-339.
2. EOS (2005): Egyptian Standards, Minced Meat Mixed With Soybean Protein. E.S: 2097 - 2005, ICS:67.120.10.
3. Belloque, J., Garcia, M.C., Torre, M. and Marina, M.L. (2002). Analysis of Soyabean Proteins in Meat Products: A Review. *Critical Reviews in Food Science and Nutrition*, 42, 5: 507-532.
4. Dyankova, S., Dyakova, A., Miteva, D. and Garbacheva, M. (2009). Study on antibacterial activity of some spice extracts used in the meat industry. *Agricultural Science and Technology*, Pilot ed., 1: 43-47.
5. Vazgecer, B., Ulu, H. and Öztan, A. (2004). Antioxidant and antimicrobial activity of spices in meat and meat products. In: Turkey 8th Food Congress, Uludağ University, Faculty of Agr., Dept. of Food Engineering, Bursa, Turkey, pp. 9.

6. Agaoglu, S., Dostbil, N. and Alemdar, S. (2007). Antimicrobial activity of some spices used in the meat industry. *Bulletin of the Veterinary Institute Pulawy*, 51: 53-57.
7. Tainter, D.R. and Grenis, A.T. (1995): Spices and Seasonings. A Food Technology Handbook. VCH, Germany: 116-119.
8. Chong-Hae, H., Ewen, T. and Gyung-Jin, B. (2008): Aerobic plate counts as a measure of hazard analysis critical control point effectiveness in a pork processing plant. *Journal of Food Protection*, 71, 6: 1248-1252.
9. Nissen, H., Maugesten, T. and Lea, P. (2001): Survival and growth of *Escherichia coli* 0157:H7, *Yersinia enterocolitica* and *Salmonella enteritidis* on decontaminated and untreated meat. *Meat Science*, 57: 291-298.
10. Deak, T. (1991): Foodborne Yeasts. *Advances in Applied Microbiology*, 36, 179-278.
11. Fleet, G. (1992). Spoilage yeasts. *Critical Reviews in Biotechnology*, 12: 1.
12. Mossel, D.A., Corry, J.E., Struijk, C.B. and Baird, R.M. (1995). Essentials of the Microbiology of Foods: A Textbook of Advanced Studies. John Wiley & Sons, Ltd, NY, USA.
13. Ahmed, A.M. and Abd El-Rahman, H.A. (2002). Effect of Ground Sage on Some Microbial Characteristics of Refrigerated Minced Beef. *Journal of King Saud University, Agric. Sci.*, 15, 1: 47-52.
14. Mitic-Ćulafic, D., Vukovic-Gacic, B., Knezevic-Vukcevic, J., Stankovic, S. and Simic, D. (2005). Comparative study on the antibacterial activity of volatiles from sage (*Salvia officinalis*L.) *Archives of Biological Science*, Belgrade, 57, 3: 173-178.
15. Oral, N., Vatansever, L., Guven, A. and Gulmez, M. (2008). Antibacterial activity of some Turkish plant hydrosols. *Kafkas. Üniv. Vet. Fak. Derg.*, 14, 2: 205-209.
16. Celikel, N. and Kavas, G. (2008). Antimicrobial properties of some essential oils against some pathogenic microorganisms. *Czech Journal of Food Science*, 26: 174–181.
17. Pokorny, J., Nguyen, H. and Korczak, J. (2008). Antioxidant activities of rosemary and sage extracts in sunflower oil. *Die Nahrung*, 41: 176-177.
18. Abdel-Hamied, A.A., Nassar, A.G. and El-Badry, N. (2009). Investigations on antioxidant and antibacterial activities of some natural extracts. *World Journal of Dairy Science*, 4, 1: 1-7.
19. Grzybowski, R., Duszkiewicz-Reinhard, W, Bugajewska, A. and Karkowska, K. (1990). Effect of a Sage Extract on Selected Microorganisms. *Przemysł Spożywczy*, 44, 11: 286-289.
20. Downes, F.P. and Ito, K. (2001). Compendium of methods for the microbiological examination of foods. 4th ed., American Public Health Association Press, Washington, D.C., USA.

21. Holt, J.G., Krieg, N.R., Sneath, P.H.A., Staley, J.T. and Williams, S.T. (1994). Bergey's manual of determinative bacteriology. 9th ed., Lippincott Williams & Wilkins, Baltimore, MD., USA.
22. Farag, R., Daw, Z., Hewedi, F. and El-Baroty, G. (1989). Antibacterial Activity of some Egyptian Spices and Essential Oils. *Journal of Food Protection*, 52, 9: 665-667.
23. Schmitz, S., Weidenboerner, M. and Kunz, B. (1993). Herbs and Spices as Selective Inhibitory of Mould Growth. *Chemie Mikrobiologie Technologie der Lebensmittel*, 15, 175-177.
24. Pitt, J.I. and Hocking, A.D.(1997). Fungi and Food Spoilage Book. 3rd Ed., Academic Press, New York, London, Toronto, Montreal, Tokyo.
25. Viljoen, B.C. and Greyling, T. (1995). Yeasts associated with cheddar and Gouda making. *International Journal of Food Microbiology*, 28:79-88.
26. Wolter, H., Laing, E. and Viljoen, B. (2000). Isolation and Identification of Yeasts Associated with Intermediate Moisture Meats. *Food Technology and Biotechnology*, 38, 1: 69-75.
27. Liu, Z. and Nakano, H. (1996). Antibacterial activity of spice extracts against food related bacteria. *Journal of the Faculty of Applied Biological Science*, Hiroshima University, 35: 181-190.
28. Hammer, K.A., Carson, C.F. and Riley, T.V. (1999). Antimicrobial Activity of Essential Oils and other Plant Extracts. *Journal of Applied Microbiology*, 86, 6: 985-990.