

Research Article

## **Optimization and kinetics of pectinase production using *Aspergillus awamori* by solid-state fermentation**

**Suresh Baladhandayutham \* and Viruthagiri Thangavelu**

Annamalai University, Annamalai Nagar – 608 002 Tamilnadu, India.

\*Email: [sure22chem@yahoo.com](mailto:sure22chem@yahoo.com)

---

### **Abstract**

Pectinase enzymes find extensive application in food and beverage industries. Pectinase production was studied in a solid-state fermentation process using rice bran and sugarcane bagasse as substrates. Optimization of media and fermentation conditions for maximum production of pectinase was carried out by one at a time procedure. Various combinations of substrates were tried to achieve maximum pectinase production. The mixed substrates consisting of 85% rice bran and 15% sugarcane bagasse gave maximum pectinase yield of 103.33U/ml during the fermentation period of 72 hours. The optimum temperature was found to be 35°C and optimum pH was found to be 5. The kinetics of pectinase production by solid state fermentation using *Aspergillus awamori* was studied. The kinetic parameters values were found to be  $K_m = 79.3$  and  $V_{max} = 1.724$  hr.ml/U .

**Keywords:** pectinolytic enzymes, fruit juice, agro-industrial wastes, fungi, India.

---

### **Introduction**

Pectinolytic enzymes catalyzing the degradation of pectic substances are of great industrial importance [1]. It has found extensive application in fruit juice industries in order to improve fruit juice yield and clarity [2]. The use of liquefying enzymes for mash treatment results in improvement of juice flow, leading to a shorter press-time, without the necessity for pressing aids [3]. At the same time, pectin is broken down to such an extent that the viscosity of mash is reduced [4]. Other areas of application include the pulp and paper industry [5], animal feed [6], retting of flax and other vegetable fibres [7], haze removal from wines [8], coffee and tea fermentation [9], oil extraction [10], purification of plant viruses [11], bio-scouring of cotton fibres [12], degumming of plant based fibres [13], protoplast fusion technology [14], textile industry [8] and waste management [14].

Among the processes used for enzyme production, solid-state fermentation (SSF) is an attractive one because it presents higher productivity per reactor volume, lower capital and operating costs, lower space requirements, simpler equipment and easier downstream processing compared to that of submerged fermentation (SmF) [15]. In addition, it permits the use of agricultural and agro-industrial residues as substrates which are converted into bulk chemicals and fine products with high commercial value such as alcohol, organic acids, fats, proteins, enzymes, etc. [16, 17]. On the other hand, there is evidence that some enzymes are less affected by catabolic repression, are more thermo stable and their optimum temperature values are higher than those obtained by SmF [18].

Major problems in the exploitation of commercial enzymes are their yield, stability, specificity and the cost of production. New enzymes for use in commercial applications with desirable biochemical and physio-chemical characteristics and a low cost of production have been the focus of research. Application of agro-industrial wastes as carbon sources in enzyme production reduces the cost of production and also helps in solving disposal problems [19, 20]. Literature highlighting the optimization, biochemical, characterization, genetics and strain improvement for pectinase production from fungi are available. However, kinetics studies on solid-state fermentative production of pectinase are lacking. Considering the biotechnological importance of fungi in enzyme engineering, the present research reported the kinetics and optimization of media and process parameters by classical method in solid state fermentation for maximum production under optimum conditions.

## **Materials and Methods**

### ***Substrate preparation***

Rice bran and sugarcane bagasse samples were obtained from an agricultural field, Salem District, Tamilnadu. The sample was made into 100 mesh (0.15 mm) fine powder by the use of laboratory grinder at 3000 rpm and was preserved in a sealed plastic bag at 4°C to prevent any possible degradation or spoilage.

### ***Microorganism and culture conditions***

Fungal strain *Aspergillus awamori* MTCC 548 was obtained from Microbial Type Culture Collection (MTCC), Institute of Microbial Technology (IMTECH), Chandigarh, India. Culture was maintained on Czapek's Agar medium. After three days incubation at 30°C the agar slants were stored at 4°C. The liquid medium for the growth of inoculum for fungi was yeast extract – malt extract – peptone – glucose medium (YMP) composed of 3 g/l of yeast extract, 3 g/l of malt extract, 5 g/l of peptone and 10 g/l of glucose.

Inocula were grown aerobically in 250 ml Erlenmeyer flasks containing the above mentioned medium at 30°C in an Environmental Shaker (Remi Scientific) at 200 rpm for 24 h. Active cells were centrifuged in a clinical centrifuge (1200 rpm), washed with sterile water and were used as inoculum. Fermentations for pectinase production were conducted on a shaker at 200 rpm in 250 ml flasks and the samples were withdrawn periodically for the analysis of enzyme concentration.

### ***Enzyme extraction***

The crude pectinase was extracted by mixing 10g of fermented materials with distilled water, stirred for 20 minutes in the shaker, filtered and then centrifuged for 20 minutes. The supernatant was used as the crude enzyme and then studied for enzymatic measurements by DNS method [21].

**Total pectinase assay**

A suitably diluted sample of 0.5ml was added to a solution containing 2 ml of 1% citrus pectin in acetate buffer (pH 4.8) in a test tube. Samples were kept at 45°C for 30 minutes in a water bath, cooled, added with 2.5 ml of DNS reagent, seethed for 5 min. Finally the contents were cooled and 10 ml of distilled water was added and was measured at 540 nm using UV/Vis Biospectrophotometer (ELICO BL 198). The concentration of  $\beta$ -galacturonic acid was determined from the standard  $\beta$ -galacturonic calibration curve.

**Solid State Fermentation (SSF)**

The powdered rice bran and sugarcane bagasse samples of different compositions were weighed (10 g/flask) and distributed into 250 ml Erlenmeyer flasks with the addition of Czapek's nutrient medium (without carbon source) to a desired solid-liquid ratio (up to 20% solid) and 0.1 M Potassium phosphate buffer (pH = 6.0), followed by sterilization for 15 min at 15 psi (121°C) in an autoclave. To the production medium  $10^8$  spores of *A. awamori* were inoculated aseptically and the flasks were then covered with cotton to allow CO<sub>2</sub> produced during fermentation to escape. The flasks were incubated in a rotary shaker (200 rpm) at 30°C for 144 h. Samples were withdrawn periodically (24h interval) and were analyzed for total pectinase enzyme activity.

**Results and Discussion****Pectinase production using different media**

The effect of substrate concentration on enzyme activity in solid-state fermentation using *Aspergillus awamori* was studied. Experiments were conducted with five different media, M<sub>1</sub>, M<sub>2</sub>, M<sub>3</sub>, M<sub>4</sub> and M<sub>5</sub> whose composition were described in Table 1. The samples were drawn at regular intervals of 24 hours and analyzed for pectinase activity. The results are given in Table 2 and Figure 1.

**Table 1. Media Composition.**

Media Composition	Medium M <sub>1</sub>	Medium M <sub>2</sub>	Medium M <sub>3</sub>	Medium M <sub>4</sub>	Medium M <sub>5</sub>
Rice Bran (g)	10	9.5	9	8.5	8
Sugar cane bagasse (g)	0	0.5	1	1.5	2
Nutrient solution (ml)	30	30	30	30	30


The pectinase enzyme activity was found to increase exponentially and reached a maximum at the end of 72 hours. A maximum of 88.65 U/ml of pectinase activity was obtained with mixed substrate of 85% of rice bran and 15% of sugarcane bagasse (Medium M<sub>4</sub>) and later on it was decreased gradually until the end of fermentation. Similar trends were observed for other mediums M<sub>1</sub>, M<sub>2</sub>, M<sub>3</sub> and M<sub>5</sub>.

The medium M<sub>4</sub> gave a maximum total pectinase activity of 88.65 U/ml and was chosen as the best medium for maximum pectinase enzyme production. The total pectinase activity for medium M<sub>4</sub> was found to be more throughout the fermentation when compared to all other media, M<sub>1</sub>, M<sub>2</sub>, M<sub>3</sub> and M<sub>5</sub>. This may be due to the addition of fibrous material as sugarcane bagasse by 15% (w/w)

increased the inter-particle spacing, possibly increasing the aeration and diffusion of nutrients and enzymes and hence resulted in the higher yield of pectinase enzyme [22]. The addition of sugarcane bagasse by 20% (w/w) decreased the yield of pectinase enzyme indicating that the microorganism was not able to hydrolyze enough cellulose and hemicellulose fibres to support mycelium formation. The use of sugarcane bagasse as support was suitable for the growth of filamentous fungi and it allowed the utilization of high concentration of substrate solutions [23].

**Table 2. Effect of Substrate Concentration on Production of Total Pectinase.**

Time h	Pectinase Activity (U/ml)				
	Medium M1	Medium M2	Medium M3	Medium M4	Medium M5
24	30.33	33.45	35.67	38.75	25.33
48	58.33	63.36	69.25	74.25	38.37
72	67.37	73.86	82.33	88.65	50.25
96	36.33	43.26	54.25	58.75	30.85
120	30	34.22	39.45	45.25	23.75



**Figure 1. Effect of Substrate Concentration on Production of Total Pectinase.**


***Effect of temperature on production of total pectinase***

The effect temperature on the pectinase enzyme production was studied using M<sub>4</sub> medium (mixed substrate with 85% rice bran and 15% sugarcane bagasse) by conducting experiments at different temperatures namely 30, 35 and 40°C by keeping all other conditions constant for the fermentation period of 120 hours. The results are given in Table 3 and Figure 2. As temperature increases from 30°C to 35°C the pectinase enzyme activity was found to increase and maximum pectinase activity of 91.67 U/ml was found at 35°C. Further increase in temperature beyond 35°C decreased the

pectinase activity until the end of fermentation. Hence optimum temperature was chosen as 35°C and was used for further studies. The decrease in enzyme activity at higher temperature may be due to enzyme denaturation.

**Table 3. Effect of Temperature on Pectinase Activity.**

Time h	Pectinase Activity (U/ml)		
	30°C	35°C	40°C
24	36.67	58.33	30.50
48	70.12	81.67	55.25
72	83.33	91.67	70
96	55.75	63.33	43.50
120	40.50	50.12	30.33




**Figure 2. Effect of Temperature on Pectinase Activity.**

#### ***Effect of initial pH on production of total pectinase***

The effect of initial pH on the pectinase production using was studied by conducting experiments at different pH namely pH3, pH4, pH5 and pH6 by keeping the temperature at 35°C and using mixed substrate with 85% rice bran and 15% sugarcane bagasse (M<sub>4</sub> medium). The results are given in Table 4 and Figure 3. As initial pH was increased from pH3 to pH5, the pectinase activity was found to increase. Further increase in initial pH beyond pH5, the pectinase activity was found to decrease. The decrease in enzyme activity at higher pH may be due to preference of fungi *A. awamori* to lower pH for its growth and metabolism. A maximum pectinase activity of 103.33 U/ml was obtained with mixed substrate for a fermentation period of 72 hours at temperature 35°C and at pH value of 5. Hence optimum pH value was chosen as pH5.

**Table 4. Effect of pH on Pectinase Activity.**


Time h	Enzyme Activity U/ml			
	pH3	pH4	pH5	pH6
24	58.33	61.67	65.25	58.33
48	81.67	83.33	86.67	75.12
72	91.67	95.50	103.33	90.25
96	63.33	65.12	76.67	55.50
120	50	56.67	63.33	38.33

**Figure 3. Effect of pH on Pectinase Activity.*****Kinetics of pectinase production***

The kinetics of pectinase production using *A. awamori* with M<sub>4</sub> medium (85% rice bran and 15% sugarcane bagasse) under optimum conditions of temperature and pH was studied. The results are given in Table 5 and Figure 4. The pectinase activity was found to increase with respect to fermentation time as shown in Figure 4 and reaches a maximum total pectinase activity of 103.33U/ml at the end of 72 hours and later on it was found to decrease until the end of the fermentation. The rate of product formation (dp/dt) was found to increase gradually and reached maximum at the end of 96 hours and later on it was found to decrease due to non-availability of substrates (as indicated in Table 5). The results showed that the maximum rate of 4.57 was obtained at the end of 72 hours and was found to be the optimum fermentation period.

**Table 5. Effect of fermentation time on pectinase production.**

Time (hrs)	Pectinase Activity(U/ml)	Rate of product formation(U/ml.hr)
0	0	0
24	65.25	1.63
48	86.67	2.95
72	103.33	4.57
96	76.67	2.45
120	63.33	1.45

**Figure 4. Effect of fermentation time on pectinase production.**


The kinetic parameters were evaluated using Lineweaver-Burk plot. The kinetic parameters were found to be  $V_{\max} = 1.724 \text{ hr.ml/U}$  and  $k_m = 79.3$ . Therefore Michaelis-Menten Kinetic model was found to be

$$V = \frac{1.724 \times S}{79.3 + S}$$

Figure 5 shows that the kinetic model was well fitted with the experimental values.

**Table 6. Comparison Between Experimental and Predicted Values.**

Substrate (rice bran) concentration (%)	Experimental values	Predicted values
100	0.924	0.961
95	0.930	0.939
90	0.937	0.916
85	0.947	0.891
80	0.883	0.865

**Figure 5. Comparison between experimental and predicted values.**

## Conclusion

Optimization of media and process parameters, namely temperature and pH, were carried out by classical method using *A. awamori* microorganism in solid state fermentation. The maximum total pectinase activity of 103.33 U/ml was achieved using the medium M<sub>4</sub> with 85% (by weight) rice bran and 15% (by weight) sugarcane bagasse. The optimum temperature was found to be 35°C and optimum pH was found to be pH5 for maximum production of pectinase. The kinetics of pectinase production was studied using medium M<sub>4</sub> under optimum conditions of temperature and pH. The kinetic parameters were determined using Lineweaver –Burk Plot with the values of  $k_m = 79.3$  and  $V_{max} = 1.724$  hr.ml/U.

### Acknowledgement

The authors express their sincere thanks to the Department of Technology, Annamalai University, for providing the necessary facilities for the successful completion of this research work.

### References

1. Alkorta, I., Garbisu, C., Llama, J.M. and Serra, J.L. (1998). Industrial applications of pectic enzymes: a review. *Process Biochemistry*, 33: 21–28.
2. Alkorta, I., Garbisu, C., Llama, J.M. and Serra, J.L. (1982). Enzymes and their uses in the processed apple industry: review. *Process Biochemistry*, 17: 35–41.
3. Charley, V.L.S. (1969). Advances in food processing using pectic and other enzymes. *Chemistry and Industry*, 20: 635- 41.
4. Mutlu, M., Srioglu, K., Demir, N., Ercan, M.T. and Acar, J. (1991). The use of commercial pectinase in fruit juice industry. Part I: Viscometric determination of enzyme activity. *Journal of Food Engineering*, 41:147-150.
5. Reid, I. and Ricard, M. (2000). Pectinase in papermaking: solving retention problems in mechanical pulps bleached with hydrogen peroxide. *Enzyme and Microbial Technolnology*, 26: 115-123.
6. Barreto de Menezes, T.J., Salva, J.G., Baldini, V.L., Papini, R.S. and Sales, A.M. (1989). Protein enrichment of citrus wastes by solid substrate fermentation. *Process Biochemistry*, 23: 167-171.
7. Hoondal, G.S., Tiwari, R.P., Tiwari, R., Dahiya, N. and Beg, Q.K. (2000). Microbial alkaline pectinases and their applications: a review. *Applied Microbiology and Biotechnology*, 59: 409–18.
8. Revilla, I. and Gonzalez-San Jose, M.L. (2003). Addition of pectolytic enzymes: an enological practice which improves the chromaticity and stability of red wines. *International Journal of Food Science and Technology*, 38:29–36.
9. Gar, J.G. (1985). Tea, coffee and cocoa. In: Wood, B.J.B., editor. *Microbiology of Fermented Foods*, Vol.2. London: Elsevier Science Ltd. 133-154.
10. Scott, D. (1978). Enzymes, industrial. In: Grayson, M., Ekarth, D. and Othmer, K. editors. *Encyclopedia of Chemical Technology*. New York: Wiley; 173–224.
11. Salazar, L. and Jayasinghe, U. (1999). Fundamentals of purification of plant viruses. In: *Techniques in plant, virology, CIP., Training Manual, J.O., Virus Purification*, International Potato Centre, Peru, 1–10.

12. Ranveer Singh, Jayani, Shivalika, Saxena and Reena, Gupta (2005). Microbial pectinolytic enzymes: a review. *Process Biochemistry*, 40: 2931-2944.
13. Kapoor, M., Beg, Q.K., Bhushan, B., Singh, K., Dadich, K.S. and Hoondal, G.S. (2001). Application of alkaline and thermostable polygalacturonase from *Bacillus* sp. MG-cp-2 in degumming of ramie (*Boehmeria nivea*) and sunn hemp (*Crotolaria juncia*) based fibres. *Process Biochemistry*, 36: 803-7.
14. Kashyap, D.R., Vohra, P.K., Chopra, S. and Tewari, R. (2001). Applications of pectinases in the commercial sector: a review. *Bioresource Technology*, 77: 215-27.
15. Acuña-Arguëlles, M.E., Gutiérrez-Rojas, M., Viniegra-González, G. and Favela-Torres, E. (1995). Production and properties of three pectinolytic activities produced by *Aspergillus niger* in submerged and solid-state fermentation. *Applied Microbiology and Biotechnology*, 43:808-814.
16. Manonmani, H.K. and Sreekantiah, K.R. (1987). Studies on the conversion of cellulase hydrolysate into citric acid by *Aspergillus niger*. *Process Biochemistry*, 22:92-94.
17. Spagnuolo, M., Crecchio, C., Pizzigallo, M.D.R. and Ruggiero, P. (1997). Synergistic effects of cellulolytic and pectinolytic enzymes in degrading sugar beet pulp. *Bioresource Technology*, 60:215-222.
18. Lozano, P., Manjón, A., Romojaro, F. and Iborra, J.L. (1988). Properties of pectolytic enzymes covalently bound to nylon for apricot juice clarification. *Process Biochemistry*, 23: 75-78.
19. Pandey, A., Soccol, C.R., Nigam, P. and Soccol, V.T. (2000). Biotechnological potential of agro-industrial residues. I: sugar cane bagasse. *Bioresource Technology*, 74, 69-80.
20. Baladhandayutham, Suresh and Thagavelu, Viruthagiri. (2010). Optimization and kinetics of pectinase enzyme using *Aspergillus niger* by solid-state fermentation. *Indian Journal of Science and Technology*, 3,743-746.
21. Miller, G.L. (1959). Use of dinitrosalicylic acid reagent for determination of reducing sugars. *Analytical Chemistry*, 31, 426-428.
22. Mitchell, D.A., Do, D.D., Greenfield, P.F. and Doell, H.W. (1991). A semi-mechanistic mathematical model for growth of *Rhizopus oligosporus* in a model solid-state fermentation system. *Biotechnology and Bioengineering*, 38, 353-362.
23. Ooijkaas, L., Weber, F.J., Buitelaar, R.M., Tramper, J. and Rinzema, A. (2000). A defined media and inert supports: their potential as solid-state fermentation production systems. *Trends in Biotechnology*, 18, 356-360.