

Polycyclic Aromatic Hydrocarbons and *n*-alkanes in Suspended Particulate Matter and Sediments from the Langat River, Peninsular Malaysia

Alireza Riyahi Bakhtiari ^a, Mohamad Pauzi Zakaria ^a, Mohammad Ismail Yaziz ^b, Mohamad Nordin Hj Lajis ^c and X. Bi ^d

 ^a Environmental Forensics Laboratory, Faculty of Environmental Studies, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia
^b Faculty of Environmental Studies, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia
^c Laboratory of Natural Products, Institute of Bioscience, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

^d State Key Laboratory of Organic Geochemistry, Guangdong Key Laboratory of Environment and Resources, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou, 510640, P.R. China

Abstract

The Langat River basin has seen rapid developments in industrialization, urbanization and dramatic population increases during the past two decades. The composition and sources of polycyclic aromatic hydrocarbons (PAHs) and aliphatic hydrocarbon (n-alkanes) concentrations were determined in surface sediments (SS) and suspended particulate matter (SPM) collected from six locations in the Langat River. The total n-alkanes concentrations (Σ HC) ranged from 5900 to 23000 µg/g in SPM and 1700 to 8600 µg/g in SS samples. Total PAHs concentrations varied from 306 to 7968 ng/g in SPM and 558 to 980 ng/g in SS. PAHs and n-alkanes were dominated by higher molecular weight compounds in SS and low-medium molecular weight compounds in SPM. Carbon preference index (CPI) values for n-alkanes in ranges C_{25-33} , C_{15-35} and C_{25-35} varied from 0.95 to 2.49 in SS and close to unity in SPM. The CPIs values indicated multiple n-alkanes sources (petrogenic and natural). PAHs isomer pairs ratios indicated multiple (petrogenic and pyrogenic) with predominance of pyrogenic PAH sources. Analysis of the possible source of PAHs and n-alkanes indicated a complicated, combined PAHs and n-alkanes source in the Langat River.

Keywords: CPI; perylene; PAHs isomer pairs ratios; suspended particular matter; surface sediment

1. Introduction

The physico-chemical properties of the PAHs largely determine their environmental behavior. Low molecular weight (LMW) PAHs, containing two or three fused rings, are more water soluble and volatile, and hence more available, than high molecular weight (HMW) PAHs containing >3 fused rings, as compared to HMW PAHs, makes them more susceptible to various degradation processes (US EPA Report, 2003).

Due to the high hydrophobicity and thereby the low water solubility, the HMW PAHs have a high affinity for organic particulate matter. With increasing the molecular weight of PAH, the tendency to sorbs onto dissolved organic matter (DOM) also increases (McCarthy and Jiminez, 1985). Therefore, the water solubility of PAH can be enhanced due to DOM and thereby co-transport of PAH with DOM occurs in the water column, especially for HMW PAHs (Maxin and Kögel-Knabner, 1995). Only lower molecular weight hydrocarbons with less than 2 and 3 condensed rings are capable of environmental transport in aqueous solution in the absence of DOM (Guggenberger *et al.*, 1996).

The Langat River is a complex ecosystem that runs through different land-uses near the city of Kuala Lumpur. Within the study area, there are seven towns located along the river. However, most of the industrial areas are located in Kajang and Bangi. There is also a rubber mill located south of Hulu Langat town and palm oil refineries near Banting.

The objectives of this study were to assess the compositional patterns and sources of PAHs and n-alkanes concentrations in surface sediment samples (SS) and suspended particulate matters (SPM) in the Langat River.

2. Materials and Methods

2.1. Sampling location

Ten surface sediments (SS) and suspended particulate materials (SPM) were collected from



Figure 1. Map of the sampling sites in the Langat River

locations in the Langat River in the west coast of Peninsular Malaysia in January 2007 (Fig. 1 and Table 1). The sediment samples were collected using an Ekman Dredge sampler. The sediment cake was placed on a stainless steel pan and the top 0-5 cm layer was taken using a precleaned stainless steel scoop. The sediment samples were then transported on ice to the laboratory and stored at -35°C until further analysis. The water was collected from the surface and middle-depth with a Niskin bottle. The water samples were then transported on ice to the laboratory. The SPM was immediately

collected after filtration of 1 to 3 L aliquots, through 0.45 Millipore μ m (47 mm diameter) preweighted filters, using the classic vacuum system and stored at -35°C until further analysis.

2.2. Chemical analysis

2.2.1. Analytical procedure for PAHs and nalkanes

The samples were purified and fractionated by a method described elsewhere (Zakaria *et al.*, 2002). Briefly, a soxhlet extractor was used for the

Table 1. Location of sampling locations and general characteristics of the suspended particulate matter and the surface sediment samples from the Langat River (19/01/07)

| Stations | | Type of sample | GPS location | | Conductivity | Salinity | Turbidity |
|------------|----------|-----------------------------------|--------------|-------------|--------------|----------|-----------|
| | Sediment | Suspended Particulate Material | | | (µs/cm) | (ppt) | (NTU) |
| S 1 | 8117 | 1 | 47N 0809449 | utm 0331379 | 35.7 | 00.0 | 168 |
| S2 | - | 1 | 47N 0806738 | utm 0322931 | 94.0 | 00.0 | 259 |
| S 3 | 1 | | 47N 0798085 | utm 0315854 | 72.6 | 00.0 | 655 |
| S4 | 1 | 1 | 47N 0769700 | utm 0316905 | 21.3 | 12.7 | 102 |
| S5 | 1 | 1 | 47N 0768258 | utm 0314266 | 28.8 | 17.6 | 164 |
| S 6 | 1 | 1 | 47N 0767248 | utm 0309944 | 44.0 | 28.2 | 25 |

extraction of lipid from the samples using 270 ml of distilled dichloromethane for 10 hours. 100 µl of the PAH surrogate internal injection standard mixture (10 ppm each component, naphthalene-d_s, anthracene- d_{10} , chrysene- d_{12} and perylene- d_{12}) was added for quality control of PAH analyses. The extracts were transferred onto the top of a 5% H₂O deactivated silica gel column. Hydrocarbons ranging from n-alkanes to PAHs with five rings were eluted with 20 ml of dichloromethane/hexane (1:3, v/v). The eluent was further fractionated using fully activated silica gel column chromatography to obtain an alkane fraction and a PAH fraction. Alkanes were eluted with 4 ml of hexane and subsequently PAHs were eluted with dichloromethane/hexane (1:3, v/v).

The PAH fraction was evaporated to approximately 1 ml transferred to a 1.5 ml amber and evaporated to dryness under a gentle stream of nitrogen and re-dissolved in 100 µl of isooctane containing *p*-terphenyl- d_{14} as an internal injection standard (IISTD) for PAHs analysis. Each n-alkane fraction was evaporated to approximately 1 ml, transferred to a 1.5 ml amber ampoule and evaporated to dryness under a gentle stream of nitrogen and re-dissolved in 100 µl of isooctane. PAHs and n-alkanes analyses were conducted using an Agilent technology 5973A quadrupole mass spectrometer integrated with an Agilent 6890 gas chromatograph. A 30 m fused silica capillary column of 0.25 mm i.d. and 0.25 µm film thickness using helium as carrier gas was used in the analysis.

GC-MS operating conditions were 70eV ionization potential with the source at 200°C and electron multiplier voltage at ~2000 eV. The injection port was maintained at 310°C and the sample was injected with splitless mode followed by purge 1 min after the injection. Column temperature was held at 70°C for 2 min, then programmed at 30°C/min to 150°C, and then 4°C/min to 310°C and held for 10 min. A selected ion monitoring method was employed after a delay of 3 min. PAHs were monitored at m/z = 128 (naphthalene), m/z = 184(dibenzothiophene), m/z = 178 (phenanthrene, anthracene), m/z = 192 (2-methylphenanthrene, 1methylphenanthrene, 2-methylanthracene), m/z =202 (fluoranthene, pyrene), m/z = 216 (1methylpyrene), m/z = 228 (chrysene, benzo[a]anthracene), m/z = 252(benzo[k]fluoranthene, benzo[e]acephenanthrene, benzo[e]pyrene, benzo[a]pyrene, perylene), m/z =278 (dibenzo[a,h]anthracene).

Each n-alkane fraction was evaporated to approximately 1 ml, transferred to 1.5 ml amber ampoule, and evaporated to dryness under a gentle

stream of nitrogen and re-dissolved in 100 µl of isooctane. It was sealed by Teflon sheet and Teflon tape, after sonication, volatile organic carbons (VOCs) are formed in the head space on the sample in the vial. n-alkane analyses were made using an Agilent technologies 5973A quadrupole mass spectrometer integrated with an Agilent 6890 gas chromatograph. A 30 m fused silica capillary column of 0.25 mm i.d. and 0.25 µm film thickness with helium as carrier gas was used in the analysis. GC-MS operating conditions were 70eV ionization potential with the source at 200°C and electron multiplier voltage at ~2000 eV. The injection port was maintained at 310°C and the sample was injected in splitless mode followed by purge 1 min after the injection. Column temperature was held at 70°C for 1 min, then programmed at 30°C/min to 150°C, 5°C/ min to 310°C and held for 10 min. A selected ion monitoring method was employed after a delay of 3 min. n-alkanes were monitored at m/z = 142 (C₁₀), $m/z = 170 (C_{12}), m/z = 198 (C_{14}), m/z = 226 (C_{16}),$ $\begin{array}{l} m/z = 254 \ ({\rm C}_{18}^{12}), \ m/z = 282 \ ({\rm C}_{20}^{10}), \ m/z = 310 \ ({\rm C}_{22}^{12}), \\ m/z = 338 \ ({\rm C}_{24}), \ m/z = 366 \ ({\rm C}_{26}), \ m/z = 394 \ ({\rm C}_{28}), \\ m/z = 422 \ ({\rm C}_{30}), \ m/z = 450 \ ({\rm C}_{32}), \ m/z = 478 \ ({\rm C}_{34}), \end{array}$ $m/z = 506 (C_{36}), m/z = 534 (C_{38}), m/z = 562 (C_{40}).$

3. Results and Discussion

3.1. n-alkanes concentrations of the Langat River

The concentrations and compositional patterns of n-alkanes and related parameters are presented in Table 2. The total n-alkanes concentrations (Σ HC) ranged from 5900 to 23000 µg/g in SPM and 1700 to 8600 µg/g in SS samples. The n-alkanes compositional patterns exhibit differences in SPM and SS samples. In the SS samples, n-alkanes are characterized by a higher abundance of long chain homologues $(nC_{27}, nC_{29}, nC_{31} \text{ and } nC_{33})$ and are dominated by nC_{29} and nC_{31} . On the other hand, in the SPM samples, n-alkanes distributions are characterized by predominance of medium molecular weights $(nC_{23}, nC_{24}, nC_{25} \text{ and } nC_{26})$ and are dominated by nC_{24} and nC_{26} . These results are consistent with those of other studies and suggest that the difference between SPM and SS samples may be attributed to easy degradation of short chain n-alkanes relative to long chain n-alkanes occurred in sediments (Meyers et al., 1984; Gagosian and Peltzer, 1986; Prahl et al., 1980; Jeng and Huh, 2004). Comparison of n-alkanes concentrations in SS samples indicated that upstream received more n-alkanes concentrations than downstream. This result may be explained by the fact that there are

| | Stations | | | | | | | | | |
|--------------------------|-----------|-----------|------------|------|-------|------|-------|------|------|--|
| | S1 | S2 | S 3 | S | 4 | S | 5 | S | 6 | |
| Compounds | SPM | SPM | SS | SS | SPM | SS | SPM | SS | SPM | |
| <i>n</i> C ₁₅ | N.D | N.D | 40 | N.D | N.D | N.D | N.D | N.D | N.D | |
| <i>n</i> C ₁₆ | N.D | N.D | 125 | N.D | N.D | N.D | N.D | N.D | N.D | |
| <i>n</i> C ₁₇ | 118 | 81 | 419 | N.D | N.D | N.D | 95 | 47 | N.D | |
| Pristane | 306 | 409 | 750 | 40 | 202 | 24 | 306 | 100 | N.D | |
| <i>n</i> C ₁₈ | 362 | 575 | 464 | 40 | 355 | 14 | 454 | 96 | N.D | |
| Phytane | 360 | 520 | 495 | 50 | 340 | 37 | 460 | 85 | N.D | |
| <i>n</i> C ₁₉ | 225 | 497 | 465 | 37 | 345 | 12 | 461 | 66 | N.D | |
| <i>n</i> C ₂₀ | 240 | 555 | 525 | 44 | 395 | 15 | 414 | 74 | N.D | |
| nC_{21} | 437 | 876 | 498 | 69 | 613 | 20 | 738 | 102 | 96 | |
| <i>n</i> C ₂₂ | 797 | 1791 | 482 | 118 | 1227 | 44 | 1533 | 229 | 189 | |
| <i>n</i> C ₂₃ | 1266 | 2690 | 549 | 172 | 1747 | 64 | 2366 | 374 | 348 | |
| <i>n</i> C ₂₄ | 1468 | 3055 | 626 | 190 | 1951 | 89 | 3048 | 461 | 550 | |
| <i>n</i> C ₂₅ | 1345 | 2088 | 870 | 248 | 1469 | 133 | 2035 | 721 | 676 | |
| <i>n</i> C ₂₆ | 1508 | 1459 | 1087 | 185 | 1036 | 140 | 1748 | 723 | 773 | |
| <i>n</i> C ₂₇ | 1044 | 1354 | 1010 | 313 | 715 | 145 | 1890 | 1095 | 667 | |
| <i>n</i> C ₂₈ | 1026 | 1019 | 1012 | 205 | 747 | 166 | 1771 | 801 | 634 | |
| <i>n</i> C ₂₉ | 1314 | 1162 | 1619 | 485 | 536 | 230 | 1995 | 1209 | 669 | |
| <i>n</i> C ₃₀ | 641 | 931 | 886 | 181 | 549 | 128 | 1664 | 550 | 572 | |
| <i>n</i> C ₃₁ | 1049 | 888 | 1732 | 627 | 570 | 264 | 1637 | 1029 | 497 | |
| <i>n</i> C ₃₂ | 807 | N.D | 828 | 172 | N.D | 115 | 1164 | 428 | 286 | |
| <i>n</i> C ₃₃ | N.D | N.D | 1182 | 301 | N.D | 148 | N.D | 570 | N.D | |
| <i>n</i> C ₃₄ | N.D | N.D | 379 | N.D | N.D | N.D | N.D | N.D | N.D | |
| <i>n</i> C ₃₅ | N.D | N.D | 625 | N.D | N.D | N.D | N.D | N.D | N.D | |
| <i>n</i> C ₃₆ | N.D | N.D | 236 | N.D | N.D | N.D | N.D | N.D | N.D | |
| Σ ΗC | 13285 | 19021 | 15704 | 3387 | 12256 | 1788 | 23013 | 8576 | 5957 | |
| CPI 25-33 | 1.03 | 1.23 | 1.47 | 2.39 | 1.09 | 1.56 | 1.00 | 1.70 | 1.00 | |
| Overall CPI 15-35 | 0.99 | 1.03 | 1.34 | 1.99 | 0.96 | 1.43 | 0.95 | 1.55 | 0.98 | |
| High CPI 25-35 | 1.02 | 1.30 | 1.48 | 2.49 | 1.10 | 1.56 | 1.03 | 1.70 | 0.96 | |
| LHC/ Σ HC | 25.0 | 17.9 | 35.3 | 51.0 | 14.9 | 44.0 | 24.0 | 45.5 | 30.8 | |

Table 2. Hydrocarbon concentrations ($\mu g/g$) and related parameters in the suspended particulate matters and the surface sediments of the Langat River

 Σ HC = Sum of $nC_{15+...+} nC_{36}$ include pristine and phytane.

SPM = Suspended particulate matter, SS = Surface Sediment.

TAR_{HC} = terrigenous/aquatic ratio = $(nC_{27}+nC_{29}+nC_{31}) / (nC_{15}+nC_{17}+nC_{19})$.

CPI ₂₅₋₃₃ = carbon preference index = $\frac{1}{2} \times [(nC_{25}+nC_{27}+nC_{29}+nC_{31}+nC_{33}-nC_{24}+nC_{26}+nC_{28}+nC_{30}+nC_{32}) + nC_{32}+nC_{32}+nC_{32}+nC_{33}+nC_{3$

 $(nC_{25}+nC_{27}+nC_{29}+nC_{31}+nC_{33} \div nC_{26}+nC_{28}+nC_{30}+nC_{32}+nC_{34})]$

Overall CPI ₁₅₋₃₅ = $(\Sigma \text{ Odds } C_{15-33} + \Sigma \text{ Odds } C_{17-35}) / 2(\Sigma \text{ Evens } C_{16-34})$

High CPI ₂₅₋₃₅ = $(\Sigma \text{ Odds } C_{25-33} + \Sigma \text{ Odds } C_{27-35}) / 2(\Sigma \text{ Evens } C_{26-34})$

LHC / Σ HC = $(nC_{27} + nC_{29} + nC_{31} + nC_{33}) / \Sigma$ HC. N.D = non detected

more human and industrial activities in the upstream (e.g. Kajang and Bangi) relative to downstream of the Langat River.

3.1.1. Carbon Preference Indices (CPI) of nalkanes

CPI is used to indicate the degree of diagenesis of straight-chain geolipids and is a numerical representation of how much of the original biological chain length specificity is preserved in geological lipids (Meyers and Ishiwatari, 1995). It is useful to determine the degree of biogenic versus petrogenic input (Mazurek and Simoneit, 1984).

In this study, the CPI values were calculated using three equations which were proposed by Bray and Evans (1961) and Cooper and Bray (1963) and listed in Table 2. The CPI values of n-alkanes in Langat River ranged from 1.00 to 2.39 in CPI ranges C_{25-33} , 0.95 to 1.55 in Overall CPI ranges C_{15-35} and 1.02 to 2.49 in High CPI ranges C_{25-35} . The CPI₂₅₋₃₃, Overall CPI₁₅₋₃₅ and High CPI₂₅₋₃₅ ranged from 1.34 to 2.49 in the SS samples relative



Figure 2. Source identification for n-alkanes using High CPI₂₅₋₃₅ (HCPI) and overall CPI₁₅₋₃₅ in Langat River sediments

the SPM samples which were close to unity (Fig. 2). The results indicate multiple sources for n-alkanes (petrogenic and natural).

3.1.2. Unresolved Complex Mixture (UCM)

Total ion chromatograms of n-alkanes showed an unresolved complex mixture (UCM) in all samples at lower carbon number maximum. The UCM concentrations were not calculated but their presence suggests the presence of degraded petroleum hydrocarbons (Simoneit, 1982; Matsumoto, 1983). It is believed that the UCM contains a mixture of alicyclic compounds (Gough and Rowland, 1990; Killops and Al-Juboori, 1990) and has a well-known linkage to degraded or weathered petroleum residues (Readman *et al.*, 1987; Venkatesan *et al.*, 1980).

3.2. PAHs composition

The concentrations of 18 PAHs, naphthalene, perylene and diagnostic ratios are presented in Table 3. Total concentration of 18 PAHs, expressed as Σ 18PAH ranged from 306 to 7968 ng/g in SS and SPM of the Langat River. The higher concentrations of total PAHs were found in upstream compared to downstream of the Langat River. This can be attributed to the fact that most of the industrial areas are located in Kajang and Bangi. In addition, there is also a rubber mill and other small and medium scale industrials located south of Hulu Langat town (Zakaria and Azril Mahat, 2006). The activities of downstream are closely related to existing port and jetties which cater to berthing and mooring facilities for various types of ships, vessels and boats. PAHs distributions were dominated by 2, 3 and 4 rings compounds in SPM samples and by 5 rings compounds in SS samples, except for Station 6. These findings of the current study are consistent with those of Luo et al. (2006) who found predominance of 5, 6-ring compounds in sediments and 2, 3-ring compounds in SPM samples from the Pearl River estuary and adjacent coastal areas in South China. Different PAHs compositional patterns between SPM and SS were also found in other regions, such as Da Ya Bay (Zhou and Maskaoui, 2003), Brighton Marina of United Kingdom (King et al., 2004), and the Baltic Sea (Witt, 1995). This may be explained to the fact that the surface water receives direct PAHs inputs, such as shipping, oil spill, dry and wet atmospheric deposition, air-water exchange etc (Luo et al., 2006). Another possible explanation for this might be due to degradation of lower molecular weight PAHs during river water column transport and after accumulation. This is due to the fact that high molecular weight PAHs are more resistant to degradation processes occurred in sediments (Cerniglia, 1992).

Naphthalene was found in abundance in SPM samples compared to SS. The range of naphthalene concentrations were from 1405 to 2420 ng/g in SPM and 410 to 440 ng/g in SS samples (Table 3). It is believed that naphthalene is a major compound in woody material (Wilcke *et al.*, 2000) and it can be supplied in large amounts to the Rivers during heavy rains in tropical environments. Naphthalene is also a product of the combustion of biomass and fossil fuel (Krauss *et al.*, 2005). It is likely therefore that the naphthalene originated from both terrigenous and pyrogenic sources. Different naphthalene compositional patterns between SPM and SS

| | Stations | | | | | | | |
|------------------|-----------|------|-----------|------|-----|------------|------|------|
| | S1 | S2 | S3 | S | 4 | S 5 | S | 66 |
| Compounds | SPM | SPM | SS | SPM | SS | SS | SPM | SS |
| Nap | 2420 | 2349 | 429 | 2348 | 438 | 417 | 1406 | 434 |
| DBT | N.D | N.D | N.D | N.D | N.D | N.D | 7 | N.D |
| Phe | 1805 | 309 | 66 | 196 | 25 | 41 | N.D | 29 |
| Ant | 352 | 437 | 22 | 215 | 15 | 32 | N.D | 46 |
| 3MPhe | 391 | 44 | 29 | 29 | 3 | 9 | 4 | 5 |
| 2MPhe | 429 | 88 | 32 | 38 | 6 | 16 | 4 | 9 |
| 2MAnt | N.D | N.D | 4 | N.D | N.D | N.D | N.D | N.D |
| 9MPhe | 133 | 70 | 40 | 22 | 6 | 12 | 4 | 5 |
| 1MPhe | 360 | 61 | 24 | 26 | 4 | 15 | 6 | 6 |
| Fluo | 813 | 136 | 59 | 95 | 24 | 55 | 27 | 43 |
| Pyr | 2381 | 166 | 169 | 84 | 63 | 138 | 43 | 81 |
| 1MPyr | 307 | 34 | 43 | N.D | 13 | 38 | N.D | 3 |
| Chry | 148 | N.D | 24 | N.D | 14 | 3 | N.D | 89 |
| BaAnt | 46 | N.D | N.D | N.D | 59 | 18 | N.D | 168 |
| BkFluo | 67 | 39 | 14 | 17 | 79 | 14 | 140 | 25 |
| BeAcep | 248 | 24 | 142 | 9 | 16 | 9 | 72 | N.D |
| BePyr | 396 | 21 | 282 | 15 | 200 | 350 | N.D | 240 |
| BaPyr | 94 | 10 | 31 | 10 | 33 | 57 | N.D | 33 |
| Per | 1.7 | 0.7 | 0.2 | 0.4 | 1.4 | 0.1 | 1.7 | 0.2 |
| DBahAnt | N.D | N.D | N.D | N.D | N.D | N.D | N.D | N.D |
| Σ18 PAHs | 7970 | 1439 | 981 | 756 | 559 | 807 | 308 | 782 |
| LMW/HMW | 1.2 | 3.9 | 0.4 | 4.6 | 0.2 | 0.3 | 0.2 | 0.2 |
| MP/P | 0.7 | 0.9 | 1.9 | 0.6 | 0.7 | 1.3 | N.A | 0.9 |
| PHE/ANT | 5.1 | 0.7 | 3.0 | 0.9 | 1.7 | 1.3 | N.A | 0.6 |
| FLT/(FLT+PYR) | 0.3 | 0.45 | 0.3 | 0.5 | 0.3 | 0.3 | 0.4 | 0.4 |
| BaA/(BaA+CHR) | 0.2 | N.A | N.A | N.A | 0.8 | 0.9 | N.A | 0.7 |
| PERY/PAIs % | 0.2 | 0.8 | 0.03 | 0.8 | 0.4 | 0.02 | 0.8 | 0.07 |
| PERY/ 2 18PAHs % | 0.02 | 0.05 | 0.02 | 0.05 | 0.3 | 0.01 | 0.6 | 0.03 |
| PYR/PERY | 1420 | 227 | 1125 | 216 | 45 | 1493 | 25 | 410 |

Table 3. PAHs concentration (ng/g) and related parameters in the surface sediments and the suspended particulate matters from the Langat River

Σ18 PAHs = Sum of (Dibenzothiophene, Phenanthrene, Anthracene, 3-Methylphenanthrene, 2-Methylphenantrene, 2-Methylphenantrene, 9-Methylphenanthrene, 1-Methylphenantrene Fluoranthene, Pyrene, 1-Methylpyrene, Chrysene, Benzo[a]Anthracene, Benzo[k]Fluoranthene, Benzo[e]Acephenanthrene, Benzo[e]Pyrene, Benzo[a]Pyrene, Dibenzo[a-h]Anthracene).

LMW/HMW = (Nap + DBT + Phe + Ant + 3Mphe + 2Mphe + 2MAnt + 9MPhe + 1MPhe + Flu)/(Pyr + 1MPyr + Chr + BaAnt + BkFluo + BeAcep + BePyr + BaPyr + Per+DBahAnt).

SPM = Suspended particulate matter.

- SS = Surface sediment.
- MP/P = (3MPhe+2MPhe+9MPhe+1MPhe)/Phe.
- PHE/ANT = Phenanthrene /Anthracene.
- FLT/(FLT+PYR) = Fluoranthene/(Fluoranthene+Pyrene).
- BaA/(BaA+CHR) = Benzo[a]Anthracene/(Benzo[a]Anthracene+Chrysene).
- PERY/PAIs = perylene/(Benzo[k]Fluoranthene+Benzo[e]Acephenanthrene+Benzo[e]Pyrene+ Benzo[a]Pyrene+perylene). PYR/PERY = Pyrene/perylene.
- $PERY / \Sigma 18PAHs = Pervlene / \Sigma 18PAHs.$
- N.A = Not applicable.
- N.D = non detected

samples can be caused by the difference in hydrophobicity between the low and high molecular weight PAHs. Low molecular weight PAHs are less hydrophobic relative to high molecular weight PAHs and therefore will be more abundant in the water column, leading to a greater proportion in suspended solids (Luo *et al.*, 2006).

Lower molecular weight versus higher molecular weight (LMW/HMW) ratios ranged from

1.16 to 4.61 in SPM and 0.17 to 0.39 in SS samples (Table 2), suggesting higher concentrations of lower molecular weight of PAHs in SPM than those of SS samples, except of Station 6 where it was 0.201 in SPM samples. Our finding are consistent with those of Luo *et al.* (2006) who found The relative content of 2, 3-ring PAHs in samples from the outer part of the estuary (on average 58.4% for samples of April 2003 and 68.6% for samples of July 2002)



Figure 3. Comparison of the MP/P ratios with a previous study on the suspended particulate matter (SPM) and the surface sediment (SS) from the Langat River

was higher than that in the inner part of the estuary (on average 46.3% for samples of April 2003 and 53.0% for samples of July 2002). This can be explain in this way that high salinity of water and low water velocity may have caused more degradation of lower molecular weight of PAHs in the estuary of the Langat River (Table 1). It may be assumed that water column processes are important events dictating the behavior and fate of PAHs in this estuary. Further investigations are required in this field.

3.2.1. Sources of PAHs in sediments and SPM

PAHs isomer pairs ratios were calculated to infer possible sources of PAHs (Table 3). In contrast to petrogenic sources, pyrogenic sources are characterized by low ratios of phenanthrene/ anthracene (PHE/ANT <15 and PHE/ANT<10) and sum of methylephenanthrenes/phenanthrene (MP/ P<1) in association with higher ratios of fluoranthene/(fluoranthene+pyrene) (FLT/ FLT+PYR)>0.4), benzo(a) anthracene/ (Benzo(a)anthracene+chrysene) (BaA/BaA+CHR) >0.35) (Budzinsky *et al.*, 1997; Benner *et al.*, 1989, 1990; Yunker *et al.*, 2002).

The MP/P ratios calculated for all SS and SPM samples were less than 2 (from 0.73 to 1.88). The results show that the PAHs in the sediments had mixed sources with predominance of pyrogenic inputs. In Fig. 3, the MP/P ratios in this study are compared to those of a previous study by Zakaria and Azril Mahat (2006). The results indicate predominance of pyrogenic inputs for PAHs in Langat River sediments at the stations except for the Stations 3 and 5.

The PHE/ANT ratios for all SS and SPM samples were less than 10 (from 0.6 to 5.1),

indicating that PAHs were mainly originated from pyrogenic sources (Fig. 4).

Based on the PAH isomer pair ratio measurements compiled by Yunker et al. (2002), FLT/FLT+PYR <0.4 implies petroleum, 0.4-0.5 implies petroleum combustion, and >0.5 implies combustion of coal, grass and wood. Ratio of BaA/ BaA+CHR <0.2 implies petroleum, from 0.2 to 0.35 indicates either petroleum or combustion, and >0.35implies combustion. The FLT/FLT+PYR ratios calculated for all SS and SPM samples were less than 0.5 (from 0.3 to 0.5). These ratios show that petrogenic input at most stations (FLT/FLT+PYR <0.4). Two out of eight samples showed values 0.4-0.5 (Stations 2 and 4), suggesting pyrogenic inputs at these two stations. The BaA/BaA+CHR ratios calculated for SS and SPM samples from four stations were less than 0.9 (from 0.2 to 0.9). The ratios FLT/FLT+PYR and BaA/BaA+CHR



Figure 4. The PHE/ANT and MP/P ratios in Langat River sediments

Table 4. Spearman's correlation coefficients (r) between Σ 18 PAHs concentrations and diagnostic ratios in surface sediments and suspended particulate matters from Langat River

| | Σ18ΡΑΗ | PHE/ANT | FLT/(FLT+PYR) | BaA/(BaA+CHR) | MP/P |
|---------------|------------------------------|----------------------------------|---|--|----------------------|
| Σ18 ΡΑΗ | 1.00 | 0.85*** | - 0.35 ^{ns} | - 0.12 ^{ns} | - 0.03 ^{ns} |
| PHE/ANT | | 1.00 | - 0.61 ^{ns} | - 0.04 ^{ns} | 0.38 ^{ns} |
| FLT/(FLT+PYR) | | | 1.00 | - 0.50 ^{ns} | - 0.47 ^{ns} |
| BaA/(BaA+CHR) | | | | 1.00 | 0.15 ^{ns} |
| MP/P | | | | | 1.00 |
| Σ18 PAHs | = Sum of (Dib 2-Methylphe | enzothiophene, enantrene, 2-M | Phenanthrene, Anthra ethylanthracene, 9-Me | acene, 3-Methylphenar thylphenanthrene, | nthrene, |

| | 1-Methylphenantrene, Fluoranthene, Pyrene, 1-Methylpyrene, Chrysene, |
|---------------|--|
| | Benzo[a]Anthracene, Benzo[k]Fluoranthene, Benzo[e]Acephenanthrene, |
| | Benzo[e]Pyrene, Benzo[a]Pyrene, Dibenzo[a-h]Anthracene). |
| PHE/ANT | = Phenanthrene / Anthracene. |
| FLT/(FLT+PYR) | = Fluoranthene / (Fluoranthene+ Pyrene). |
| BaA/(BaA+CHR) | = Benzo[a]Anthracene / (Benzo[a]Anthracene+ Chrysene). |

MP/P = Sum of (3-Methylphenanthrene, 2-Methylphenanthrene, 9-Methylphenanthrene, 1-Methylphenanthrene) / Phenanthrene.

Levels of significant: * = p < 0.05; ** = p < 0.01; *** = p < 0.001; ^{ns} = p > 0.05

indicated multiple PAHs sources (petrogenic and pyrogenic).

Correlation between $\Sigma 18$ PAHs and diagnostic ratios calculated to identify sensitivity of the PAH ratios. The results exhibit a significant positive correlation (r = 0.854, p = 0.007) between $\Sigma 18$ PAHs and PHE/ANT at the 99% confidence level, suggesting predominance of pyrogenic versus petrogenic inputs in Langat River sediments. The ratios of FLT/FLT+PYR, MP/P and BaA/ BaA+CHR were not significantly correlated with $\Sigma 18$ PAHs, suggesting they are not sensitive enough to be used to distinguish petrogenic PAHs from pyrogenic PAHs in an area like the Langat River, where pollutants come from multiple sources (Table 4).

3.2.2. Perylene

The concentrations of perylene and diagnostic ratios such as, perylene/penta-aromatic isomers (PERY/PAIs), pyrene/perylene (PYR/PERY) and perylene/total PAHs (PERY/ Σ 18PAHs) are presented in Table 3. For pyrogenic PAHs, perylene concentration is about 1-4% of total PAHs (Wakeham *et al.*, 1980a; 1980b; Wang *et al.*, 1999b). In addition, perylene concentrations relative to penta-aromatic isomers greater than 10% indicate probable diagenic inputs, whereas those less than 10% indicate a probable pyrogenic origin of the compound (Baumard *et al.*, 1998a; Readman *et al.*, 2002). The concentrations of pyrene relative to those of perylene were compared to identify different source of high temperature anthropogenic PAHs (Venkatesan 1988; Silliman *et al.*, 2001). The perylene concentrations ranged from 0.09 to 1.71 ng/g. The high ratio of PYR/PERY ranged from 25 to 1500 and low percentages of PERY/PAIs <1% and PERY/ Σ 18PAHs <1%, suggest pyrogenic input for perylene in Langat River sediments.

4. Conclusions

The results of PAHs and n-alkanes analysis of surface sediments and suspended particulate matters from the Langat River were used to assess likely sources of these contaminants. Total concentrations and compositional patterns of PAHs and n-alkanes showed differences in the SPM and the SS samples. Low molecular weight of PAHs and n-alkanes dominated in the SPM samples. The concentrations of high molecular weight of PAHs and n-alkanes were higher in SS samples than those in SPM samples. This can be attributed to the differences in hydrophobicity and degradation processes between the low and high molecular weight PAHs; it could also result in the compositional difference between SS and SPM samples.

PAHs isomer pairs ratios suggested a complicated, combined PAHs source (petrogenic and pyrogenic) with predominance of pyrogenic inputs in the study area. The perylene concentrations and related perylene ratios (PERY/PAIs, PERY/ Σ 18PAHs and PYR/PERY) point to pyrogenic

inputs for perylene in the study area. The high concentration of naphthalene in all sediment samples and higher concentrations in the SPM relative the SS samples reveal an additional source for naphthalene in Langat River sediments. This can be attributed to woody plants in this tropical environment.

The results of the CPIs values for n-alkanes in ranges C_{25-33} , C_{15-35} and C_{25-35} exhibited higher values in the SS samples relative the SPM samples, indicating mixed petrogenic and natural sources with predominance of petrogenic inputs for n-alkanes in the SPM.

Acknowledgment

This study was supported by the Ministry of Science, Technology and Innovation, Malaysia (MOSTI) who provided partial funding for this project (Science Fund Project (No. 04-01-04-SF0092).

References

- Baumard P, Budzinski H, Michon Q, Garrigues P, Burgeot T, Bellocq J. Origin and bioavailability of PAHs in the Mediterranean Sea from mussel and sediment records. Estuarine Coastal Shelf Science 1998; 47: 77-90.
- Bray EE, Evans ED. Distribution of n-paraffins as a clue to recognition of source beds. Geochimica et Cosmochimica Acta. 1961; 22: 2-15.
- Benner BA, Bryner NP, Wise SA, Mulholland GH, Lao RC, Fingas MF. Polycyclic aromatic hydrocarbons emissions from combustion of crude oil on water. Environment Science and Technology 1990; 24: 1418-27.
- Benner BA, Gordon GE, Wise SA. Mobile sources of atmospheric polycyclic aromatic hydrocarbons: a roadway tunnel study. Environment Science and Technology 1989; 23: 1269-78.
- Budzinski H, Garrigues P, Bernard G, Bellocq J, Hinrichs K, Rullkotter J. Identification of polycyclic aromatic hydrocarbons in sediments from the Amazon fan: occurrence and diagenetic evolution. *In:* Proceedings of the Ocean Drilling Program, Scientific Results (*Eds*: Flood RD, Piper DJW, Klaus A, Peterson LC). Ocean Drilling Program, College Station (TX), 1997: 555-64.
- Cerniglia CE. Biodegradation of polycyclic aromatic hydrocarbons. Biodegradation 1992; 3: 351-68.
- Cooper JE, Bray EE. A postulated role of fatty acids in petroleum formation. Geochimica et Cosmochimica Acta. 1963; 27: 1113-27.
- Gagosian RB, Peltzer ET. The importance of atmospheric input of terrestrial organic material to deep sea sediments. Organic Geochemistry 1986; 10: 661-69.
- Gough MA, Rowland J. Characterization of unresolved complex mixtures of hydrocarbons in petroleum. Nature 1990; 344: 648-50.

- Guggenberger G, Pichler M, Hartmann R, Zech W. Polycyclic aromatic hydrocarbons in different forest soils: mineral horizons. Journal of Plant Nutrition and Soil Science. 1996; 159: 565-73.
- Jeng WL, Huh CA. Lipids in suspended matter and sediments from the East China Sea Shelf. Organic Geochemistry 2004; 35: 647-60.
- Killops SD, Al-Juboori M. Characterisation of the unresolved complex mixture (UCM) in the gas chromatograms of biodegraded petroleums. Organic Geochemistry 1990; 15: 147-60.
- King AJ, Readman JW, Zhou JL. Dynamic behaviour of polycyclic aromatic hydrocarbons in Brighton marina, UK. Marine Pollution Bulletin 2004; 48: 229-39.
- Krauss M, Wilcke W, Martius C, Bandeira AG, Marcos VB, Garcia MVB, Amelung W. Atmospheric versus biological sources of polycyclic aromatic hydrocarbons (PAHs) in a tropical rain forest environment. Environment Pollution 2005; 135: 143-54.
- Luo XJ, Chen SJ, Mai BX, Yang QS, Sheng GY, Fu JM. Polycyclic aromatic hydrocarbons in suspended particulate matter and sediments from the Pearl River Estuary and adjacent coastal areas, China. Environment Pollution 2006; 139: 9-20.
- Matsumoto G. Comparative study on organic constituents in polluted and unpolluted inland aquatic environments-V. Organic carbons and hydrocarbons in sediments. Water Research 1983; 17: 823-30.
- Maxin C, Kögel-Knabner I. Partitioning of PAH to DOM: Implications on PAH mobility in soils. European Jurnal of Soil Science 1995; 46: 193-204.
- Mazurek MA, Simoneit BRT. Characterization of biogenic and petroleum-drived organic matter in aerosols over remote, rural, and urban area. *In:* Identification and Analysis of Organic Pollutants in Air (*Ed*: Keith LH). Ann Arbor Science, Woburn, MA, 1984: 353-70.
- McCarthy JF, Jiminez B. Interactions between polycyclic aromatic hydrocarbons and dissolved humic material: Binding and dissociation. Environment Science and Technology 1985; 19: 1072-76.
- Meyers PA, Ishiwatari R. Organic matter accumulation records in lake sediments. *In*: Physics and Chemistry of Lakes (Eds: Lerman A, Imboden DM, Gat JR). Berlin, Springer, 1995; 279-328.
- Meyers PA, Leenheer MJ, Eadie BJ, Maule SJ. Organic geochemistry of suspended and settling particulate matter in Lake Michigan. Geochimica et Cosmochimica Acta. 1984; 48: 443-52.
- Prahl FG, Bennett JT, Carpenter R. The early diagenesis of aliphatic hydrocarbons and organic matter in sedimentary particulates from Dabob Bay, Washington. Geochimica et Cosmochimica Acta.. 1980; 44: 1967-76.
- Readman JW, Fillmann G, Tolosa I, Bartocci J, Villeneuve P, Catinni C, Mee LD. Petroleum and PAH contamination of Black Sea. Marine Pollution Bulletin 2002; 44: 48-62.

- Readman JW, Mantoura RFC, Rhead MM. A record of polycyclic aromatic hydrocarbons (PAH) pollution obtained from accreting sediments of the Tamar estuary, UK: Evidence for nonequilibrium behaviour of PAH. Science of the Total Environment.1987; 66: 73-94.
- Simoneit BRT. Some applications of computerized GC-MS to the determination of biogenic and anthropogenic organic matter in the environment. International Journal of Environmental Analytical Chemistry. 1982; 12: 177-93.
- Silliman J, Meyers P, Eadie B, Klump J. A hypothesis for the origin of perylene based on its low abundance in sediments of Green Bay, Wisconsin. Chemical Geology 2001; 177: 309-22.
- US EPA Report, 2003. Report on bioavailability of chemical wastes with respect to potential for soil bioremediation. EPA/600/R-03/076.
- Venkatesan MI. Occurrence and possible sources of perylene in marine sediments—a review. Marine Chemistry 1988; 25: 1–27.
- Venkatesan MI, Brenner NS, Tuth E, Bonilla J, Kaplan IR. Hydrocarbons in age-dated sediment cores from two basins in the Southern California Bight. Geochim. Cosmochim. Acta, 1980; 44: 789-802.
- Wakeham SG, Achaffner C, Glger G. Polycyclic aromatic hydrocarbons in recent lake sediments: II. Compounds derived from biogenic precursors during early diagenesis. Geochim Cosmochim Acta. 1980a; 44: 415-29.
- Wakeham SG, Schaffner C, Giger W. Polycyclic aromatic hydrocarbons in recent lake sediments-I. Compounds having anthropogenic origins. Geochim. Cosmochim. Acta. 1980b; 44: 403-13.
- Wang Z, Fingas M, Shu YY, Sigouin L, landriault M, Lambert P, Turpin R, Campagna P, Mullin J. Quantitative characterization of PAHs in burn residue and soot samples and differentiation of pyrogenic PAHs from petrogenic PAHs-the 1994 mobile burn study. Environment Science and Technology 1999b; 33: 3100-09.
- Wilcke W, Amelung W, Martius C, Garcia MVB, Zech W. Biological sources of polycyclic aromatic hydrocarbons (PAHs) in the Amazonian rain forest. Journal of Plant Nutrition and Soil Science. 2000; 163: 27-30.
- Witt G. Polycyclic aromatic hydrocarbons in water and sediments of the Baltic Sea. Marine Pollution Bulletin 1995; 31: 237-48.
- Yunker MB, Macdonald RW, Vingarzan R, Mitchell RH, Goyette D, Sylvestre S. PAHs in the Fraser river basin: a critical appraisal of PAH ratios as indicators of PAH source and composition. Organic Geochemistry 2002; 33: 489-515.
- Zakaria MP, Azril Mahat A. Distribution of Polycyclic aromatic hydrocarbon (PAHs) in sediments in the Langat estuary. Costal Marine Science 2006; 30(1): 387-95.
- Zakaria MP, Takada H, Tsutsumi S, Ohno K, Yamada J, Kouno E, Kumata H. Distribution of polycyclic

aromatic hydrocarbons (PAHs) in rivers and estuaries in Malaysia: a widespread input of petrogenic PAHs. Environment Science and Technology 2002; 36: 1907-18.

Zhou JL, Maskaoui K. Distribution of polycyclic aromatic hydrocarbons in water and surface sediments from DaYa Bay, China. Environmental Pollution 2003; 121: 269-81.

Received 18 December 2008 Accepted 20 February 2009

Correspondence to

Dr. Mohamad Pauzi Zakaria Environmental Forensics Laboratory, Faculty of Environmental Studies, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia Tel: (603)8946-6738 Fax: (603)8946-7463 E-mail: mpauzi@env.upm.edu.my