
Green insecticide from herbal essential oils against house fly, *Musca domestica* L. (Muscidae: Diptera)

J. Sinthusiri^{*}, M. Soonwera and P. Boonmeesupmak

Entomology and Environment Program, Faculty of Agricultural Technology, King Mongkut's Institute of Technology Ladkrabang, Ladkrabang, Bangkok, Thailand

J. Sinthusiri, M. Soonwera and P. Boonmeesupmak (2013) Green insecticide from herbal essential oils against house fly, *Musca domestica* L. (Muscidae: Diptera). Journal of Agricultural Technology 9(6):1453-1460.

The insecticidal effect of essential oils from *Cinnamomum verum* (cinnamon), *Myristica fragrans* (nutmeg) and *Syzygium aromaticum* (clove) against house fly (*Musca domestica* L.) using WHO susceptibility test. Each was applied in ethyl alcohol at concentrations of 1%, 5% and 10% (v/v). Ten percent concentrations of *M. fragrans* and *S. aromaticum* oils were found more effective than cypermethrin (positive control). They gave 100 % knockdown at 15 min, KT_{50} values of 5.94 and 3.80 min, respectively. In addition, 10% *M. fragrans* and *S. aromaticum* oils gave 98% and 100% mortality rate at 24 hrs and LC_{50} were 2.81 and 1.21%, respectively. While, cypermethrin showed 100% knockdown at 30 min, KT_{50} value of 7.28 min and LC_{50} value of 8.45%. The data demonstrated the potential of nutmeg and clove oils to be used as green insecticide against house fly.

Keywords: insecticidal, herbal essential oil, house fly, *Musca domestica* L.

Introduction

The house fly, *Musca domestica* L. (Diptera, Muscidae) is one of the most synanthropic insects in the world. It imposes itself on humans and eats all what is available, food and wastes. Therefore, housefly is considered mechanical vectors of pathogens (bacteria, protozoa and viruses) to human and causes economic problems to all farm animals (cattle, pig and poultry) (Olsen *et al.*, 2001; Sangmaneeet *et al.*, 2005; Jesikha, 2012). These pathogens may cause many contagious diseases such as food poisoning, diarrhea, cholera, typhoid and paratyphoid, shigellosis, and anthrax (Banjo *et al.*, 2005; Fasanella *et al.*, 2005; Yap *et al.*, 2008). They are also vectors of a variety of eggs from worm parasites (Wattanachai *et al.*, 1996; Ugbogu *et al.*, 2006). Moreover, myiasis in humans have been reported (Dogra and Mahajan, 2010).

^{*} Corresponding author: Jirisuda Sinthusiri, e-mail: jiri_ja@yahoo.com

House fly control is largely based upon the use of chemical insecticides. Majority of the chemical insecticides are harmful to man and animals, some of which are not easily degradable and spreading toxic effect. The increased use of these insecticides may enter into the food chain, and thereby, the liver, kidney, etc., may be irreversibly damaged (Dhanasekaran *et al.*, 2013). Currently, the use of eco-friendly, biodegradable insecticides from plants to control insects are gaining importance and botanicals have been found to be effective. Essential oils from plants have been reported to have potential insecticidal effects adult house flies, such as the oils from *Cymbopogon citratus*, *Mentha piperita* and *Lavandula angustifolia* showed 100% knockdown at 30 and 60 min., and LC₅₀ values were 2.22, 2.62 and 3.26%, respectively (Sinthusiri and Soonwera, 2013). The essential oils from *Minthostachys verticillata*, *Hedeoma multiflora*, *Citrus sinensis*, *Citrus aurantium*, *Eucalyptus cinerea* and *Artemisia annua* showed LC₅₀ of 0.5, 1.3, 3.9, 4.8, 5.5 and 6.5 mg/dm³ in 30 min, respectively (Palacios *et al.*, 2009a; Palacios *et al.*, 2009b). Tarelli *et al.*, (2009) found KT₅₀ at 2.3 and 19.0 min. for monoterpenes of eucalyptus oil (eucalyptol) and peppermint oil (menthone), respectively. Hence, essential oils from various plants may be potential alternatives to used house fly control. *Cinnamomum verum* J. Presl. (*Lauraceae*) *Myristica fragrans* Houtt. (*Myristicaceae*) and *Syzygium aromaticum* (L.) Merrill. & L.M. Perry (*Myrtaceae*) trees are the local plants in Asia, Southeast Asia and other parts of the tropics, their oils and volatile constituents are widely used in the prevention and treatment of human illnesses because they have the bioactivity of antibacterial, antiviral, antifungal, antioxidant, anti-inflammatory and pesticidal activity (Latha *et al.*, 2005; Bhamarapravati *et al.*, 2006; Mayaud *et al.*, 2008; Pandey and Singh, 2011; Djiani and Dicko, 2012).

Therefore, the present study aimed to investigate *Cinnamomum verum* (cinnamon), *Myristica fragrans* (nutmeg) and *Syzygium aromaticum* (clove) essential oils as an insecticidal against adult house fly.

Materials and methods

Rearing of houseflies

House fly adults were collected from Udomsuk market, Huatakae, Ladkrabang, Bangkok, Thailand. They were reared in gauzier cotton cages (30x30x30 cm) at 30-35°C and 70-80% relative humidity in the laboratory of Entomology and Environmental Programme, Plant Production Technology Section, Faculty of Agricultural Technology, King Mongkut's Institute of Technology Ladkrabang (KMITL), Bangkok, Thailand. They were fed with 10% syrup soaked in cotton wool and powder milk 30 g. Then, 300 g of

Mackerel fish were placed on a plastic tray (18x25x9 cm) lined with sterile coconut husks for house flies to feed and lay their eggs. Four days-old of adult house flies were used for the study.

Essential oils

The essential oils from *Cinnamomum verum* J. Presl. (cinnamon), *Myristica fragrans* Houtt. (nutmeg) and *Syzygium aromaticum* (L.) Merrill. & L.M. Perry (clove) were tested in this study. They were extracted by water distillation at the laboratory of KMITL and prepared as 1%, 5%, and 10% solutions in ethyl alcohol (v/v). All formulations were kept at room temperature before testing.

Chemical Insecticide

Cypermethrin (Kumakai[®]) was purchased from M.D Manufacturing Co. Ltd., Pranakornsri Ayutaya province, Thailand and used as standard.

Insecticide susceptibility test

The insecticide susceptibility test adapted from WHO guidelines (2006) used a susceptibility test kit. Each tube was marked with either a red spot or a green spot. Ten house flies were collected and released into the green spot tubes lined with clean filter paper (Whatman[®] No.1, 12x 15 cm) with a movable slide attached. The same size filter paper was cut for the bioassay. The filter paper was impregnated with 2 ml of each essential oil. After the filter paper dried for 20 minutes, it was inserted into the red spot tube for exposure to the house flies. The house flies were exposed to the treated paper for 1 hour in each tube. Knockdown rates were recorded at 5, 10, 30 and 60 minutes. At the end of the exposure period, house flies were released back into the green spot tube and provided with 10% syrup. Mortality was observed after 24 hours. Each test was performed in 5 replicated with simultaneous control sets; the positive controls were impregnated with cypermethrin and the negative controls were ethyl alcohols.

Statistical analysis

The data was pooled and analyzed by standard probit analysis to obtain KT_{50} and LC_{50} . The knockdown and mortality data were statistically analyzed using one-way ANOVA and the data means were compared by Duncan's multiple range test. Significant difference was considered at $P < 0.05$. The

susceptibility results were determined for each insecticide using WHO criteria (WHO, 1998): 98-100% mortality indicated susceptibility, 80-97% mortality suggested possible resistance needing confirmation and mortality <80% suggested resistance. If mortality exceeded 20% in the control, the whole test should be rejected and repeated. If mortality in the controls were between 5-20%, results with the treated samples should be corrected using Abbott's formula.

Results

Percent knockdown and KT_{50} values of three essential oils at 1%, 5% and 10% concentrations against house flies were shown in Table 1. At 1% concentration, the essential oil from *S. aromaticum* had high knockdown against house flies, there were 90% knockdown at 60 min, KT_{50} was 16 min. While cypermethrin (positive control) gave KT_{50} at 21.62 min. At 5% concentration, *M. fragrans*, *S. aromaticum* and *C. verum* showed 96% knockdown at 60 min, KT_{50} were 12.87, 12.91 and 18.17 min, respectively. Cypermethrin showed 94% knockdown at 60 min, KT_{50} was 15.90 min. At 10% concentration, *S. aromaticum* and *M. fragrans* had 100% knockdown at 15 min, KT_{50} were 3.80 and 5.94 min, respectively. Meanwhile, cypermethrin showed 100% knockdown at 30 min, KT_{50} was 7.28 min. Ethyl alcohols (negative control) percent knockdown could not be found.

The mortality rate, susceptibility and LC_{50} values of house flies for three essential oils were shown in Table 2. At 1% concentration, house flies were resistant to all essential oils and cypermethrin with the mortality rates ranging from 16 to 40%. At 5% concentration, house flies were susceptible (S) to *S. aromaticum* with 100% mortality rates. They were resistance susceptible (RS) to *M. fragrans* and *C. verum*, with the mortality rates of 92 and 82%, respectively. Cypermethrin gave 28% mortality rate which showed resistance to the house flies. At 10% concentration *M. fragrans* increased the mortality rates to 98%. Thus, house flies were susceptible (S) to *S. aromaticum* and *M. fragrans*, resistance susceptible (RS) to *C. verum*, and resistance (R) to cypermethrin. There were significant differences of mortality rates which calculated by using one-way ANOVA. All essential oils gave LC_{50} values at 24 hours after exposure less than cypermethrin. LC_{50} values of *S. aromaticum*, *M. fragrans* and *C. verum* were 1.21, 2.81 and 3.07%, respectively while cypermethrin was 8.45%.

Table 1. KT_{50} values and percent knockdown of house flies by 1, 5 and 10 % concentration of three essential oils after exposure at 5, 15, 30 and 60 minutes

Herbal essential oils	1% Concentration				KT_{50}		5% Concentration				KT_{50}		10% Concentration				KT_{50}	
	5 min	15 min	30 min	60 min	min	n	5 min	15 min	30 min	60 min	min	n	5 min	15 min	30 min	60 min	min	n
<i>C. verum</i>	0	0 ^C	20±1 0.00 B	38±25. 90 ^B	65 .7 3	0 ^A	58±1 1.00 ^B	92±8 .40 ^A	96±8. 90 ^A	18 .1 7	6±8. 90 ^C	86± 7.10 B	94± 5.50 B	96±5. 50 ^B	14 .5 3			
<i>M. fragrans</i>	0	0 ^C	16±2 0.70 B	52±17. 90 ^B	57 .2 1	0 ^A	90±1 2.20 ^A	96±5 .50 ^A	96±8. 90 ^A	12 .8 7	32±3 0.30 ^B	100 A	100 A	100 ^A	5. 94			
<i>S. aromaticum</i>	0	78±4 .50 ^A	82±4 .50 ^A	90±7.1 0 ^A	16 .0 0	0 ^A	84±8 .90 ^A	84±8 .90 ^A	96±5. 50 ^A	12 .9 1	90±1 0.00 ^A	100 A	100 A	100 ^A	3. 80			
cypermethrin	0	54±1 1.40 B	84±1 1.40 A	90±10. 00 ^A	21 .6 2	2±4 .50 A	70±1 8.70 ^A	92±1 3.00 A	94±8. 90 ^A	15 .9 0	18±2 1.70 ^B	96± 5.50 A	100 A	100 ^A	7. 28			
ethyl alcohol	0	0 ^C	0 ^C	0 ^C	0	0 ^A	0 ^D	0 ^D	0 ^B	0	0 ^C	0 ^C	0 ^C	0 ^C	0			
CV%	N A	30.3 7	32.4 6	30.64		547 .72	20.59	12.9	8.35		68.51	4.96	2.71	2.7				

KT_{50} , 50% knockdown time: NA, not available

Mean % knockdown in each column followed by the same letter are not significantly different (one-way ANOVA and Duncan's multiple range test)

Table 2. The mortality rates, susceptibility and LC_{50} values of house flies to three essential oils

Herbal essential oils	1% concentration		5% concentration		10% concentration		LC_{50} at 24 hrs
	%Mortality	Susceptibility	%Mortality	Susceptibility	%Mortality	Susceptibility	
<i>C. verum</i>	24±19.50 ^{AB}	R	82±11.00 ^B	RS	88±8.37 ^B	RS	3.07
<i>M. fragrans</i>	16±26.10 ^{BC}	R	92±8.40 ^{AB}	RS	98±4.47 ^A	S	2.81
<i>S. aromaticum</i>	40±14.10 ^A	R	100±0 ^A	S	100±0 ^A	S	1.21
cypermethrin	24±0.89 ^{AB}	R	28±1.92 ^C	R	60±1.22 ^C	R	8.45
ethyl alcohol	0 ^C	R	0 ^D	R	0 ^D	R	NA
CV%	79.29		14.41		8.5		

LC_{50} , 50% lethal concentration: Mean % mortality in each column followed by the same letter are not significantly different (one-way ANOVA and Duncan's multiple range test).

NA, not available: S, Susceptible is defined as 98-100% mortality; RS, Resistance susceptible is defined as 80-97% mortality; R, Resistance is defined as <80% mortality

Discussions

Insecticidal effect of *C. verum*, *M. fragrans* and *S. aromaticum* oils against four days old adult house flies. The results showed that 10% *M. fragrans* and *S. aromaticum* oils had high knockdown rates and susceptibility to houseflies more than cypermethrin. Similarly, Palacios *et al.* (2009a) reported *M. fragrans* oil had effect to house flies, with LC_{50} value was 8.8 mg/dm³. Jung *et al.* (2007) reported *M. fragrans* seed compound which (1R)-(+)-camphor, (1S)-

(-)-camphor, dipentene, (1R)-(+)- β -pinene, and (+)- α -terpineol were more toxic to adult female german cockroach than propoxur. Phuakbuakhao and Soonwera (2010) found *S. aromaticum* oil had highly effected to american cockroach adult which 100% mortality at 5 min and LT_{50} was 0.59 min. Omara *et al.* (2013) found *S. aromaticum* oil gave highly significant effect against american cockroach nymphs and adult by fumigant test, with LC_{50} value of 1.06 μ l and 8.20 μ l and 100% mortality at a concentration of 7.5 and 17.5 μ l/L of air, respectively. This oil was also most effective to adult mosquitoes and cowpea weevil. Makhaik *et al.* (2005) found *S. aromaticum* oil gave LC_{50} and LC_{95} values at 0.5 and 0.9%, respectively for *Culex Quinquefasciatus* and 2.7 and 5.3%, respectively for *Aedes aegypti*. Mahfuz and Khalequzzaman (2007) found *S. aromaticum* oil was highly toxic against *Callosobruchus maculates*, with LD_{50} of 92.81 and 69.62 μ g/cm² after expose at 24 and 48 hrs., respectively.

Moreover, *M. fragrans* and *S. aromaticum* has been used in medicine, flavoring food, dental products, perfumes and lotions. In East Asia, the seed of *M. fragrans* were used in medical properties for digestive and kidney problems. In the British herbal Pharmacopoeia, Myristica seed oil is indicated as specific for flatulent dyspepsia, nausea, diarrhea and dysentery (Jung *et al.*, 2007). Dried flower of *S. aromaticum* has long been considered to use in folk medicine as analgesic, toothache, sore throat, respiratory disorder, digestive system ailment, dyspepsia, gastritis, diarrhea, diuretic and carminative (Pandey and Singh, 2011; Debjit *et al.*, 2012). Both essential oils have been found inhibited against food borne pathogens such as *Staphylococcus aureus*, *Escherichia coli*, *Bacillus anthracis*, *B. mycoides*, *B. pumilus*, *B. subtili*, *Saccharomyces cervisiae*, and *Shigella* spp. (Latha *et al.*, 2005; Pandey and Singh, 2011).

In conclusion, *M. fragrans* and *S. aromaticum* oils, act not only toxic to adult house flies but also many useful to human. Thus, essential oils from *M. fragrans* and *S. aromaticum* were interesting alternative insecticidal to control house flies. Because, they safe to human, animal and environmental friendly.

Acknowledgements

The authors are highly grateful to Faculty of Agricultural Technology, King Mongkut's Institute of Technology Ladkrabang (KMITL) and The National Research Council of Thailand (NRCT) for providing financial assistance to carry out this study.

References

- Banjo, A.D., Lawal, O.A. and Adeduji, O.O. (2005). Bacteria and fungi isolated from housefly (*Musca domestica* L.) larvae. African Journal of Biotechnology 4:780-784.

- Bhamarapravati, S., Juthapruth, S., MahaChai, W. and Mahady, G. (2005). Antibacterial activity of *Boesenbergia rotunda* (L.) Mansf. and *Myristica Fragrans* Houtt. against *Helicobacter pylori*. Songklanakalin Journal of Science and Technology 28(1):157-163.
- Debjit, B., Kumar, K.P.S., Yadav, A., Srivastava, S., Paswan, S. and Dutta A.S. (2012). Recent trends in Indian traditional herbs *Syzygium aromaticum* and its health benefits. Journal of Pharmacognosy and Phytochemistry 1(1):13-23.
- Dhanasekaran, S., Krishnappa, K., Anandan, A. and Elumalai, K. (2013). Larvicidal, ovicidal and repellent activity of selected indigenous medicinal plants against malarial vector *Anopheles stephensi* (Liston.), dengue vector *Aedes aegypti* (Linn.) and *Culex tritaeniorhynchus* (Giles.) (Diptera: Culicidae). Journal of Agricultural Technology 9(1):29-47.
- Djiani, A. and Dicko, A. (2012). The therapeutic benefits of essential oils. Available from URL: <http://cdn.intechweb.org/pdfs/29979.pdf>. [2012/8/3]
- Dogra, S.S. and Mahajan, V.K. (2010). Oral myiasis caused by *Musca domestica* larvae in child. International Journal of Pediatric Otorhinolaryngology 5:105-107.
- Fasanella, A., Scasciamacchia, S., Garofolo, G., Giangaspero, A., Tarsitano, E. and Adone, R. (2005). Evaluation of the house fly *Musca domestica* as a mechanical vector for an anthrax. Plos one 5:1-5.
- Jesikha, M. (2012). Evaluation of larvicidal efficacy of *Aloe Vera* extract against *Musca domestica*. IOSR Journal of Environmental Science, Toxicology and Food Technology 2(2):1-3.
- Jung, W.C., Jang, Y.S., Hieu, T.T., Lee, C.K. and Ahn, Y.J. (2007). Toxicity of *Myristica fragrans* Seed Compounds Against *Blattella germanica* (Diptera: Blattellidae). Journal of Medical Entomology 44(3):524-529.
- Latha, P.G., Sindhu, P.G., Suja, S.R., Geetha, B.S., Pushpangadan, P. and Rajasekharan, S. (2005). Pharmacology and chemistry of *Myristica Fragrans* Houtt. Journal of Spices and Aromatic Crops 14(2):94-101.
- Mahfuz, I. and Khalequzzaman, M. (2007). Contact and fumigant toxicity of essential oils against *Callosobruchus maculatus*. University Journal of Zoology, Rajshahi University 26:63-66.
- Makhaik, M., Naik, S.N. and Tewary, D.K. (2005). Evaluation of anti-mosquito properties of essential oils. Journal of Scientific and Industrial Research 64:129-133.
- Mayaud, L., Carricajo, A., Zhiri, A. and Aubert, G. (2008). Comparison of bacteriostatic and bactericidal activity of 13 essential oils against strains with varying sensitivity to antibiotics. Letters in Applied Microbiol 47(3):167-173.
- Olsen, A.R., Gecan, J.S., Ziobro, G.C. and Bryce, J.R. (2001). Regulatory action criteria for filth and other extraneous materials V. Strategy for evaluating hazardous and nonhazardous filth. Regulatory Toxicology and Pharmacology 33:363-392.
- Omara, S.M., Al-Ghamdi, K.M., Mahmoud, M.A.M. and Sharawi, S.E. (2013). Repellency and fumigant toxicity of clove and sesame oils against American cockroach (*Periplaneta Americana* L.). African Journal of Biotechnology 12(9):963-970.
- Palacios, S.M., Bertoni, A., Rossi, Y., Santander, R. and Urzua, A. (2009a). Efficacy of essential oils from edible plants as Insecticides against the house fly, *Musca domestica* L. Molecules 14:1938-1947.
- Palacios, S.M., Bertoni, A., Rossi, Y., Santander, R. and Urzua, A. (2009b). Insecticidal activity of essential oils from native medicinal plants of central Argentina against the house fly, *Musca domestica* (L.). Parasitology Research 106:207-212.
- Pandey, A. and Singh, P. (2011). Antibacterial activity of *Syzygium aromaticum* (clove) with metal ion effect against food borne pathogens. Asian Journal of Plant Science and Research 1(2):69-80.

- Phuakbuakhao, N. and Soonwera, M. (2010). Effect of herbal essential oils to control American cockroach (*Periplaneta americana*). Proceeding of the 16th Asain Agricultural Symposium and 1st International Symposium on Agricultural Technology, Bangkok, Thailand. Scientific Publ, Bangkok, 25-27 August 2010. Tokai University and King Mongkut's Institute of Technology Ladkrabang. pp 659-662.
- Sangmaneedet, S., Kanistanon, K., Papirom, P. and Tesiri, T. (2005). Uses of Thai medicinal herb (*Derris elliptica* (Roxb.) Benth) in control of fly larvae population and its application in the treatment of cutaneous Myiasis in animals. Khon Kaen University Research Journal 10:22-30.
- Sinthusiri, J. and Soonwera, M. (2013). Efficacy of herbal essential oils as insecticides against the housefly, *Musca Domestica* L. Southeast Asian Journal of Tropical Medicine and Public Health 44(2):188-196.
- Tarelli, G., Zerba, E.N. and Alzogaray, R.A. (2009). Toxicity to vapour exposure and tropical application of essential oils and monoterpenes on *Musca domestica* (Diptera: Muscidae). Journal of Economic Entomology 102:1383-1388.
- Ugbogu, O.C., Nwachukwu, N.C. and Ogbuagu, U.N. (2006). Isolation of Salmonella and Shigella species from houseflies (*Musca domestica* L.) in Utaru, Nigeria. African Journal of Biotechnology 5:1090-1091.
- Wattanachai, P., Sooppapathom, K., Tintanon, B. and Phun-urai, P. (1995). Susceptibility of house fly (*Musca domestica*) to Different compound in Thailand, 1993-1995. Journal of Health Science 5:572-575.
- World Health Organization. (2006). Guidelines for testing mosquito adulticides for indoor residual spraying and treatment of mosquito nets. WHO/CDS/NTD/WHOPES/GCDPP/2006.3.
- World Health Organization. (1998). Test procedures for insecticide resistance monitoring in malaria vectors, bio-efficacy and persistence of insecticides on treated surfaces. WHO/CDS/CPC/MAL/98.12.
- Yap, K.L., Kalpana, M. and Lee, I.L.L. (2008). Wings of the commom house fly (*Musca domestica* L.): importance in mechanical transmission of *Vibrio cholerae*. Tropical Biomedicine 25:1-8.

(Received 26 August 2013; accepted 31 October 2013)