
Insecticidal efficacy of essential oil from *Cinnamomum zeylanicum* Blume and its two major constituents against *Callosobruchus maculatus* (F.) and *Sitophilus oryzae* (L.)

Jyotika Brari and D.R. Thakur*

Department of Biosciences, Himachal Pradesh University, Shimla, 171005, India.

Jyotika Brari and D.R. Thakur (2015) Insecticidal efficacy of essential oil from *Cinnamomum zeylanicum* Blume and its two major constituents against *Callosobruchus maculatus* (F.) and *Sitophilus oryzae* (L.). *Journal of Agricultural Technology* 11(6): 1323-1336

Essential oil isolated from *Cinnamomum zeylanicum* Blume was analyzed by mass spectroscopy (MS) and two of its main constituents were identified. Contact and fumigant activity of *C. zeylanicum* essential oil and its two constituents viz., cinnamaldehyde and linalool were examined against two stored product pests, *Callosobruchus maculatus* F. (Coleoptera: Bruchidae) and *Sitophilus oryzae* L. (Coleoptera: Curculionidae). Responses varied with respect to doses of compounds, insect species and exposure time. In contact activity assay, the cinnamon oil at 1.2 mg/cm² caused 98 and 80% mortality against *C. maculatus* and *S. oryzae* adults respectively, after 24 hours of treatment. 0.3 mg/cm² of cinnamaldehyde caused 100 % mortality in *C. maculatus* whereas, similar dose of linalool caused 89% mortality in *S. oryzae* after an exposure period of 12 hrs. The fumigant action as compared to contact toxicity indicated high percentage of mortality for adult insects even at low doses of compounds and less duration of exposure. The essential oil of cinnamon along with its major constituents demonstrates higher contact and fumigant toxicity to *C. maculatus* than *S. oryzae* adults. Further, cinnamaldehyde showed higher toxicity than linalool and it may be attributed to its chemical structure.

Keywords: Essential oil, cinnamaldehyde, linalool, contact toxicity, fumigant toxicity.

1. Introduction

Stored insect pests are a major problem throughout the world as they significantly reduce the quantity and quality of grain. The post-harvest grain losses due to insect pests and other bio-agents ranged from 10 to 40% (Raja *et al.* 2001; Papachristos and Stamopoulos 2002). It is evident from literature that the cowpea weevil, *Callosobruchus maculatus* (F.) and the rice weevil, *Sitophilus oryzae* (L.) are considered as the most widespread and destructive

Corresponding Author: jyotika58brari@gmail.com , drdr4@rediffmail.com

primary insect pests of stored legumes and cereals (Park *et al.* 2003). The main method of grain protection and to avoid or control insect infestations is the use of chemical agents since, it is the simplest and most cost effective means of dealing with stored product pests (Hidalgo *et al.* 1998). But the excessive use of traditional chemical insecticides leads to a number of serious problems, like persistence in the atmosphere, resistance to chemical insecticides, pests resurgence, exclusion of economically beneficial insects, toxicity to humans and environment and higher cost of crop production (Khan and Selman 1989). So, there is an urgent need to develop safe alternatives to conventional fumigants and insecticides to control the insect infestation in stored grain products. Continued screening for such systems of grain protection that target the pest species more accurately is required (Cox 2004). Plant derived insecticides as compared to synthetic ones are safer for the environment, generally cost effective, easy to handle and used by small industries and farmers (Belmain *et al.* 2001). Botanical pesticides are often active against a number of species, are often biodegradable, nontoxic and appropriate for use in integrated pest management (Kim *et al.* 2003). In this context, plant extracts including essential oils can play an important role in protecting stored products against insect infestations. Essential oils derived from plants are volatile in nature (Guenther 1948) and their constituents have revealed adequate activity as botanical pesticides (Singh and Upadhyah 1993). Spices are considered as rich sources of essential oils and are known to be effective against various insect pests including stored product insect pests (Jacobson 1989). Clove oil was found toxic to *S. oryzae* and castor oil to *C. maculatus* and *C. phaseoli* in stored conditions (Pacheco *et al.* 1995). Fumigant toxicity of essential oils from several spices like anise (*Pimpinella anisum* L.) and peppermint (*Mentha piperita* L.) was found against four major stored product pests including, *Ryzopertha dominica* (F.), *Tribolium castaneum* (Herbst), *S. oryzae* and *Orzyaephilus surinamensis* (L.) (Shaaya *et al.* 1991).

In this work, we have examined the insecticidal activity of essential oil from *Cinnamomum zeylanicum* and its two major components cinnamaldehyde and linalool against *C. maculatus* and *S. oryzae* adults. This insecticidal study will put emphasis on the practice of essential oils and their components as botanical insecticides against variety of insect pests.

Objectives: • To reduce the negative impacts on human health and environment natural products are the best alternative to synthetic pesticides.

• In the present study cinnamon oil and its two major constituents were shown great efficiency against the two stored products insect pests (*C. maculatus* and *S. oryzae*), responsible for post-harvest losses.

- Essential oils are target specific, biodegradable, less expensive, promising grain protectants and relatively less harmful to environment and mammalian health.

2. Materials and methods

2.1. Chemicals

Cinnamaldehyde (>99% purity) and linalool (>99% purity) were purchased from Sigma Aldrich and tested by bioassay. Methanol was purchased from SD fine-chem Ltd, Mumbai, India. The chemical structure of compounds has been shown in **figure 1**.





Figure 1. Chemical structure of linalool and cinnamaldehyde.

2.2. Insect culture

Two species of insects *S. oryzae* and *C. maculatus* were reared in the culture. Cultures of *S. oryzae* and *C. maculatus* were maintained in the laboratory without exposure to any insecticide on rice grain and cowpea respectively, in glass containers at $25 \pm 1^\circ\text{C}$ and 50 - 60% rh.

2.3. Extraction of essential oil from cinnamon

Cinnamon barks (5g) was grinded into fine powder and then transferred into a 100 ml round-bottomed flask (RBF). 20 ml of ethyl acetate along with few pieces of anti-bumping granules were added to the round-bottomed flask and then a water-cooled condenser was attached to the flask followed by refluxing of the mixture for 15-20 minutes with constant heating at 70°C . The mixture was cooled to room temperature and filtered into a clean conical flask. After that the mixture was passed through anhydrous sodium sulphate to remove the moisture. Transfer the Filtrate was transferred into the round-bottomed flask (RBF) and the solvent from mixture removed under reduced pressure in a rotary flash evaporator at 45°C and oily residue left behind is the essential oil.

2.4 Mass spectroscopy

Mass spectra of cinnamon oil was recorded on Bruker micrOTOF Q II Mass spectrometer. The prime components of essential oil were identified by mass spectroscopy and mass spectrum of essential oil isolated from cinnamon has been shown in **figure 2**.




Figure 2. Mass spectrum of cinnamon oil.

2.5. Bioassay

a) Contact toxicity

Insecticidal activities of cinnamon oil and its two constituents against adults of *C. maculatus* and *S. oryzae* have been determined by direct contact application as cited in literature (Kim *et al.* 2003; Usha Rani 2010). Cinnamon oil and its constituents were prepared in methanol at concentrations of (0.6, 0.8 & 1.2 mg/cm²) and (0.1, 0.2 & 0.3 mg/cm²) respectively. Aliquots of 1 ml of each solution were applied on filter papers (Whatman No. 1). Solvent was allowed to evaporate for 10 - 15 minutes before the introduction of insects. Then each dried paper was placed at the bottom of a Petri dish (5.5 cm × 1.2 cm) and 10 adults each of *C. maculatus* and *S. oryzae* were placed in each Petri dish and covered with a lid. Controls received only 1000 µl methanol alone. Each set of treatment was repeated three times and number of dead insects in each petri-dish was counted at an interval of 3, 6, 8, 12, 24 48 hour respectively. Percentage mortality was calculated by using the Abott formula.

b) Fumigant toxicity

Vapour toxicity of essential oil and its constituents against the adult insects were determined via impregnated paper assay following the method of Park *et al.* (2002) with some modifications. Plastic jars of 250 ml capacity with screw lids were used as exposure chambers. Different doses of 30 µl, 20 µl and 10 µl of cinnamaldehyde and linalool dissolved in methanol were applied to a circular filter paper (Whatman No. 1, 3 cm diameter). The treated filter paper discs were then introduced into the plastic jars (250 ml capacity) to achieve final concentrations of 0.04, 0.08, 0.12, 1.2, 2.4 and 3.6 µl ml⁻¹, respectively, with respect to volume of the jars. After allowing the solvent to evaporate for 10 - 15 minutes, the filter paper was attached to the inner surface of the screw lid of the jar using adhesive tape. Ten adults of each insect species (2 to 4 days old) were released into each jar containing 50 seeds. Insect mortalities were determined and calculated after 3, 5, 7, 12 and 24 hours from exposure. Three replicates were set up for each dose and control.

2.6 Statistical analysis

All the data concerning mortality were corrected by using Abbott's formula (Abbott 1925). Tests for contact and fumigant toxicity were performed in triplicate and data presented are mean \pm SE. The mean values were compared by one-way ANOVA and Tukey's multiple comparison tests using software SPSS, version 11.5.

3. Results

3.1. Components of cinnamon essential oil

The mass spectroscopy has been used to elucidate the components of the cinnamon oil. The essential oil from *C. zeylanicum* contains different component have been summarized in **table 1**.

Table 1. Chemical constituents of the essential oil (*C. zeylanicum*) and parent ion peak (m/Z) values of components.

Components	m/Z	
	calculated	Observed
alpha-pinene	136.23	136.20
Myrcene	154.24	154.20
Linalool	154.13	154.20
1,8-cineole	154.24	154.20
Cinnamaldehyde	132.05	132.34
Eugenol	164.08	164.23
beta-caryophyllene	204.36	204.24
cinnamyl acetate	176.21	176.30
para-cymene	134.21	134.10
alpha-terpineol	154.25	154.20

The parent ion peak (m/Z) for the different components of cinnamon oil has been observed in mass spectrum at m/z 132.34, 134.10, 136.20, 154.20, 164.23, 176.30 and 204.244 and these signals account for cinnamaldehyde (calculated m/z = 132.05), para-cymene (calculated m/z = 134.21), alpha-pinene (calculated m/z = 136.23), myrcene (calculated m/z = 154.24), linalool (calculated m/z = 154.13), 1,8-cineole (calculated m/z = 154.24), alpha-terpineol (calculated m/z = 154.25), eugenol (calculated m/z = 164.08), cinnamyl acetate (calculated m/z = 176.21) and beta-caryophyllene (calculated m/z = 204.36). The mass spectroscopy revealed that the *C. zeylanicum* oil

composed of a mixture of components mentioned above. A detailed survey of literature regarding the insecticidal activity of essential oils suggests that the phenolic/aromatic compounds affected the insecticidal activity positively as compared to non-aromatic compounds. Therefore in the present work one aromatic (cinnamaldehyde) and one non-aromatic (linalool) component of essential oil were chosen to explore the insecticidal activities.

3.2. Contact activity with treated filter paper

The contact activity method has been widely used to investigate the toxicity of insecticides. The findings of contact toxicity unveiled that the mode of action of essential oil and its components against adults of *C. maculatus* and *S. oryzae* was dosage and exposure time dependent. It has been noticed that the cinnamon essential oil showed 52 and 65% mortality against adult *C. maculatus* at a dose of 0.6 and 0.8 mg/cm² respectively. Further increase the concentration of oil to 1.2 mg/cm², yielded higher mortality of 98% against adult *C. maculatus*. A similar trend of mortality versus concentration has been shown by essential oil against *S. oryzae* (**table 2**) but mortality rate was slightly lower in case of *S. oryzae*. But no mortality was obtained in the control within the same time period.

Table 2. Contact activities of essential oil of cinnamon against adults of *C. maculatus* and *S. oryzae*.

Essential oil	Dose (mg/cm ²)	Mortality%	
		<i>C. maculatus</i> (24 hour*)	<i>S. oryzae</i> (48 hour*)
	0.6	52.6 ±3a	45.4 ±5a
0.8	65.3 ±2b	56.5 ±4b	
1.2	98.2±1c	95.3 ±2c	

*Exposure time and % values are mean ($n = 3$) ± SE. The means within a column followed by same letter are not significantly different from each other according to ANOVA and Tukey's comparison tests.

Results in **figure 3** displayed the components of essential oil cinnamaldehyde and linalool showed strong toxicity against both the insect species than the cinnamon oil. In a test with *C. maculatus* adults, cinnamaldehyde at a dose of 0.3 mg/cm², caused 88 and 100% mortality at the time interval of 6 and 8 hour treatment respectively. But in case of *S. oryzae* maximum mortality of 95% was achieved by cinnamaldehyde after 12 hour of treatment at a dose of 0.3 mg/cm².




Figure 3. Represents the percentage mortality of a) *C. maculatus* and b) *S. oryzae* respectively, exposed to cinnamaldehyde and linalool by contact activity assay.

However, in linalool a dose of 0.3 mg/cm² caused 66, 75 and 86% mortality in *C. maculatus* within 6, 8 and 12 hour of treatment respectively and in *S. oryzae* 52, 65 and 89% mortality within same exposure time (table 3 and 4).

Table 3. Contact activities of constituents of essential oil from cinnamon against *C. maculatus*.

	Dose (mg/cm ²)	Mortality%		
		6 hour*	8 hour*	12 hour*
Linalool	0.3	66.2 ±2cd	75.6 ±3de	86.3 ±1a
	0.2	42.5±6bc	52.4 ±5a	76.4 ±3de
	0.1	30.8±4e	51.6 ±4a	65.5 ±3cd
Cinnamaldehyde	0.3	88.2 ±2cde	100 ±0b	100 ±0b
	0.2	78.3 ±3ef	88.4 ±4d	92.4 ±1ef
	0.1	56.2 ±5d	65.6 ±2fg	86.8 ±2a

*Exposure time and % values are mean ($n = 3$) ± SE. The means within a column followed by same letter are not significantly different from each other according to ANOVA and Tukey's comparison tests

Furthermore, the minimum mortality achieved by cinnamaldehyde against *C. maculatus* and *S. oryzae* was 56 and 50% respectively, at a concentration of 0.1 mg/cm² and 6 hour of exposure. Similarly, linalool also showed mortality of 30 and 28% at lowest dose of 0.1mg/cm² against *C. maculatus* and *S. oryzae* respectively, 6 hour after treatment. The study revealed that the mortality was greatly affected by exposure time as well as concentration (table 3 and 4). The

results also shows that the toxicity was more pronounced in cinnamaldehyde than linalool and this may be accredited to aromatic nature of cinnamaldehyde.

Table 4. Contact activities of constituents of essential oil from cinnamon against *S. oryzae*.

	Dose (mg/cm ²)	Mortality%		
		6 hour*	8 hour*	12 hour*
Linalool	0.3	52.4 ±1bc	65.4 ±3a	89.1 ±1de
	0.2	39.5 ±1cd	48.2 ±1e	60.5 ±2cd
	0.1	28.2 ±4de	35.8 ±2fg	48.9 ±1cde
Cinnamaldehyde	0.3	78.3 ±2e	88.6 ±2de	95.2 ±5fg
	0.2	69.2 ±1f	72.6 ±4b	88.6 ±3e
	0.1	50.7 ±1g	65.7 ±1a	82.4 ±2c

*Exposure time and % values are mean ($n = 3$) ± SE. The means within a column followed by same letter are not significantly different from each other according to ANOVA and Tukey's comparison tests.

3.3. Fumigant toxicity

The efficacy of investigated compounds (essential oil, cinnamaldehyde and linalool) has further been evaluated by fumigation toxicity. The cinnamon oil at a dose of 1.2 μlml^{-1} caused 36 and 32% mortality of *C. maculatus* and *S. oryzae* respectively, after 7 hour of treatment whereas, at higher concentration (3.6 μlml^{-1}) the essential oil caused 100 and 95% mortality against both the species of insects at same exposure of time (**figure 4**). *C. maculatus* was found to be more susceptible to essential oil than *S. oryzae*.




Figure 4. Percentage mortality of *C. maculatus* and *S. oryzae* exposed to fumigant action of cinnamon essential oil.

At a dose of $0.12 \mu\text{ml}^{-1}$ both the components of essential oil caused 100% mortality in both insect species after 7 hour of treatment (**figure 5**). Similarly, at a concentration of $0.04 \mu\text{ml}^{-1}$, cinnamaldehyde caused 66 and 82% mortality of *C. maculatus* 3 and 5 hour after exposure whereas, 63 and 79% mortality was achieved in *S. oryzae* at the same duration of exposure respectively (**table 5**).




Figure 5. Represents the percentage mortality of a) *C. maculatus* and b) *S. oryzae* respectively, exposed to cinnamaldehyde and linalool by fumigant action.

Table 5. Fumigant toxicity of essential oil/ constituents against adults of *C. maculatus*.

	Doses (μml^{-1})	Mortality %		
		3 hour*	5 hour*	7 hour*
Cinnamon oil	3.6	52.2 ±1b	69.4 ±2cd	100 ±0a
	2.4	40.5 ±3bc	46.7 ±3b	68.5 ±5fg
	1.2	28.6 ±2c	25.8 ±2efg	36.4 ±2gh
Linalool	0.12	60.8 ±3a	67.3 ±5cd	100 ±0a
	0.08	52.3 ±1b	54.4 ±ef	88.6 ±1def
	0.04	34.4 ±2de	46.2 ±3b	72.5 ±3ghi
Cinnamaldehyde	0.12	78.4 ±4cd	89.1 ±2a	100 ±0a
	0.08	68.1 ±2cde	88.5 ±2a	100 ±0a
	0.04	66.5 ±5cde	82.2 ±4cde	100 ±0a

*Exposure time and % values are mean ($n = 3$) ± SE. The means within a column followed by same letter are not significantly different from each other according to ANOVA and Tukey's comparison tests.

However linalool at dose of 0.04 μml^{-1} gave 46 and 34% mortality of *C. maculatus* and *S. oryzae* respectively, 5 hour after treatment. Again no mortality was observed under controls. The increased exposure time and concentrations increased the mortality of insects. The fumigant activity of cinnamaldehyde against *C. maculatus* increased by 1 - 2.2 folds as concentration increases by 3 times at any exposure of time whereas, the activity of linalool increases by 1.3 - 1.7 folds against the same species of insect. Further, the fumigant action of the essential oil pronounced by 1.8 - 2.9 folds against *C. maculatus* with 3 times increase in dose concentration at any exposure of time. The vapour toxicity of cinnamaldehyde was found more than the cinnamon essential oil and linalool. Also the mortality rate was significantly affected by concentration in case of essential oil than other investigated compounds (cinnamaldehyde and linalool). *C. maculatus* was found to be more susceptible to essential oil and its constituents.

Table 6. Fumigant toxicity of essential oil/ constituents against adults of *S. oryzae*.

	Doses (μml^{-1})	Mortality %		
		3 hour*	5 hour*	7 hour*
Cinnamon oil	3.6	49.7 \pm 1ab	56.5 \pm 2cd	95.2 \pm 2b
	2.4	35.7 \pm 3bc	48.4 \pm 5de	65.2 \pm 5ef
	1.2	20.8 \pm 2bcd	23.8 \pm 2bcd	32.8 \pm 2efg
Linalool	0.12	56.3 \pm 1cd	62.5 \pm 5ef	100 \pm 0a
	0.08	42.3 \pm 4cde	48.6 \pm 4de	82.4 \pm 1e
	0.04	32.4 \pm 2a	34.3 \pm 3bc	69.9 \pm 3gh
Cinnamaldehyde	0.12	68.3 \pm 4b	86.2 \pm 2d	100 \pm 0a
	0.08	66.8 \pm 2b	85.1 \pm 1d	100 \pm 0a
	0.04	63.5 \pm 5b	79.4 \pm 4efg	96.1 \pm 3b

*Exposure time and % values are mean ($n = 3$) \pm SE. The means within a column followed by same letter are not significantly different from each other according to ANOVA and Tukey's comparison tests.

Discussions

The present study unveiled that the essential oil of *C. zeylanicum* and its two components exhibits contact and fumigant toxicity against the adults of both insect species. The toxic effects of cinnamon essential oil are attributed to its major constituent monoterpenes which are highly volatile and possess high fumigant toxicity. Many plant derived materials such as monoterpenoids have fumigant action against a variety of insect pests attributed to their high volatility (Coats *et al.* 1991; Shaaya *et al.* 1997; Ahn *et al.* 1998). Monoterpenoids (limonene, linalool, terpineol, carvacrol and myrcene) are the main insecticidal constituents of many essential oils effective against stored product insects (Regnault-Roger *et al.* 1995). Based on previous investigation (Regnault-Roger and Hamroui 1995) two monoterpenes cinnamaldehyde and linalool were selected from different constituents of cinnamon essential oil for comparative study of their contact and fumigant action with essential oil of cinnamon against two stored product pests. The essential oil of cinnamon along with its major constituents revealed higher toxicity to *C. maculatus* than adults of *S. oryzae*. El- Nahal *et al.* (1984) studied the toxic effect of *Acoras calamus* essential oil on five stored product insect species and found the declining order of susceptibility as *C. chinensis*, *S. granaries*, *S. oryzae*, *T. confusum* and *R. dominica*. Both the prime components showed better insecticidal activity than the essential oil. In the present study contact toxicity assay demonstrated highest mortality within 12 hours of treatment with cinnamaldehyde and linalool while complete mortality by fumigation was reached within 7 hours of treatment in both the insect species. However in contact paper assay cinnamon oil caused 98 and 95% mortality at a high dose of 1.2 mg/cm² against *C. maculatus* and *S. oryzae* adults respectively in a time period of 24 and 48 hours of treatment. The results indicated that the fumigant activity of cinnamon oil, linalool and cinnamaldehyde was more as compared to their contact toxicity. The insecticidal activity of mustard oil, horse radish and foeniculum fruit extract against *Lasioderma serricorne* adults was recorded more in closed cups than in open ones and also insecticidal mode of action of these materials was largely attributed to their fumigant action. (Kim and Ahn *et al.* 2003). Ahn *et al.* (1998) reported that the monoterpene carvacrol has a wide range of insecticidal activity against various agricultural, stored product and medical insect pests, and possess fumigant activity. The phenylpropenes (E)-anethole, estragole and monoterpene fenchone exhibits fumigant activity against adults of *S. oryzae*, *C. chinensis* and *L. serricorne* (Kim and Ahn 2001). The adulticidal activity of cinnamon essential oil and its constituents was found to be both dose and exposure time dependent. Cinnamaldehyde showed higher toxicity than linalool may be due to its aromatic structure. Fumigant activity and

reproductive inhibition induced by monoterpenes with chemical skeleton having a non-substituted phenolic moiety was found more against *Acanthoscelides obtectus* (Regnault–Roger and Hamraoui 1995). Present study demonstrates that essential oil and its constituent monoterpenes can be used for managing stored product insects in enclosed spaces such as storage bins, glasshouses and buildings etc. because of their fumigant action. But Karr and Coats (1992) have reported that when monoterpenoids are used as potential insecticides, direct toxicity as well as adverse effects on biotic potential must be considered in the evaluation of overall insecticidal efficacy.

It is possible that essential oils from plants and their derived constituents have sufficient potential to replace the more problematic fumigants and insecticides. Therefore they might be considered as better consolidates in the avenue of botanical insecticides for pest management. But further investigations are required to increase knowledge horizon for the effective and widespread use of these technologies especially on commercial bases.

Acknowledgement

Jyotika Brari is thankful to Himachal Pradesh University for research facilities. Jyotika Brari is also thankful to Dr. D. R. Thakur for his guidance in this research work.

References

- Abbott WS (1925) A method for computing the effectiveness of an insecticide. *J Econ Entomol* 18:265–267.
- Ahmed SM, Eapen M (1986) Vapour toxicity and repellency of some essential oils to insect pests. *Indian Perfumer* 30:273–278.
- Amos TG, Williams P, Du Guesclin PB, Schwarz M (1974) Compounds related to juvenile hormone: activity of selected terpenoids on *Tribolium castaneum* and *T. confusum*. *J Econ Entomol* 67:474–476.
- Coats JR, Karr LL, Drewes CD (1991) Toxicity and neurotoxic effects of monoterpenoids. In *Insects and Earthworms*. ACS Symposium Series. 449 (Naturally Occurring Pest Bioregister) 305-316.
- Finney DJ (1971) *Probit Analysis*, 3rd Edition. Cambridge University Press, London, UK. Grundy, D.L., Still, C.C., 1985. Inhibition of acetylcholinesterases by pulegone-1,2-epoxide. *Pestic Biochem Physiol* 23:383–388.
- Hough-Goldstein JA (1990) Antifeedant effects of common herbs on the Colorado potato beetle (Coleoptera: Chrysomelidae). *Environ Entomol* 19:234–238.

- Huang Y, Tan WMJ, Kini RM, Ho HS (1997) Toxic and antifeedant action of nutmeg oil against *Tribolium castaneum* (Herbst) and *Sitophilus zeamais* Motsch. J Stored Prod Res 33:289–298.
- Huang Y, Ho SH (1998) Toxicity and antifeedant activities of cinnamaldehyde against the grain storage insects, *Tribolium castaneum* (Herbst) and *Sitophilus zeamais* Motsch. J Stored Prod Res 34:11-17.
- Isman MB (2006) Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. Annu Rev Entomol 51: 45–66
- Karr LL, Coats JR (1988) Insecticidal properties of d-limonene. J Pestic Sci 13:287–290.
- Kim DH, Ahn YJ (2001) Contact and fumigant activities of constituents of *Foeniculum vulgare* fruit against three Coleopteran stored-product insects. Pest Manag Sci 57:301–306.
- Kordali S, Aslan I, Calmasur, O, Cakir A (2006) Toxicity of essential oils isolated from three *Artemisia* species and some of their major components to granary weevil, *Sitophilus granarius* (L.) (Coleoptera: Curculionidae). Ind Crops Prod 23:162–170.
- Lee S, Tsao R, Peterson C, Coats JR (1997) Insecticidal activity of monoterpenoids to western corn rootworm (Coleoptera: Chrysomelidae), twospotted spider mite (Acari: Tetranychidae), and house fly (Diptera: Muscidae). J Econ Entomol 90:883–892.
- Lee BH, Choi WS, Lee SE, Park BS (2001a) Fumigant toxicity of essential oils and their constituent compounds towards the rice weevil, *Sitophilus oryzae* (L.). Crop Prot. 20:317-320.
- Lee SE, Lee, BH, Choi WS, Park BS, Kim JG, Campbell BC (2001b) Fumigant toxicity of volatile natural products from Korean spices and medicinal plants towards the rice weevil, *Sitophilus oryzae* (L.). Pest Manag Sci 57:548-553.
- Lee SE, Peterson CJ, Coats JR (2003) Fumigation toxicity of monoterpenoids to several stored product insects. J Stored Prod Res 39:77–85.
- Lee BH, Annis PC, Tumaalii F, Choi, WS (2004) Fumigant toxicity of essential oils from the Myrtaceae family and 1, 8–cineol against 3 major stored–grain insects. J Stored Prod Res 40: 553– 564.
- Liu ZL, Ho HL (1999) Bioactivity of essential oils extracted from *Evodia rutaecarpa* Hook f. et Thomas against the grain storage insects, *Sitophilus zeamais* Motsch. and *Tribolium castaneum* (Herbst). J Stored Prod Res 35:317–328.
- Misra G, Pavlostathis SG (1997) Biodegradation kinetics of monoterpenes in liquid and soil-slurry systems. Appl Microbiol Biotechnol 47:572–577.

- Ogendo JO, Belmain SR, Deng AL, Walker DJ (2003) Comparison of toxic and repellent effects of *Lantana camara* L. with *Tephrosia vogelii* Hook and a synthetic pesticide against *Sitophilus zeamais* Motschulsky in maize grain storage. *Insect Sci Appl* 23:127–135.
- Prates HT, Santos JP, Waquil JM, Fabris JD, Oliveira AB, Foster JE (1998) Insecticidal activity of monoterpenes against *Rhyzopertha dominica* (F.) and *Tribolium castaneum* (Herbst). *J Stored Prod Res* 34:243–249.
- Regnault-Roger C, Hamraoui A (1993) Efficiency of plants from South of France used as traditional protectants of *Phaseolus vulgaris* L. against its bruchid *Acanthoscelides obtectus* Say. *J Stored Prod Res* 29:259 - 264.
- Regnault-Roger C, Hamraoui A (1995) Fumigant toxic activity and reproductive inhibition induced by monoterpenes on *Acanthoscelides obtectus* (Say) (Coleoptera), a bruchid of kidney bean (*Phaseolus vulgaris* L.). *J Stored Prod Res* 31:291-299.
- Rice PJ, Coats JR (1994) Insecticidal properties of several monoterpenoids to the house fly (Diptera: Muscidae), red flour beetle (Coleoptera: Tenebrionidae), and southern maize rootworm (Coleoptera: Chrysomelidae). *J Econ Entomol* 87:1172–1179.
- Stamopoulos DC (1991) Effects of four essential oil vapours on the oviposition and fecundity of *Acanthoscelides obtectus* (Say) (Coleoptera: Bruchidae): laboratory evaluation. *J Stored Prod Res* 27:199–203.
- Tripathi AK, Prajapati V, Kumar S (2003) Bioactivities of l-carvone, d-carvone, and dihydrocarvone toward three stored product beetles. *J Econ Entomol* 96:1594–1601.
- Tsao R, Lee S, Rice PJ, Jensen C, Coats JR (1995) Monoterpenoids and their synthetic derivatives as leads for new insect control agents. In: Baker, D.R., Fenyves, J.G., Basarab, G.S. (Eds.), *Synthesis and Chemistry of Agrochemicals IV*. American Chemical Society, Washington, DC, pp. 312–324.
- Watanabe K, Shono Y, Kakimizu A, Okada A, Matsuo N, Satoh A, Nishimura H (1993) New mosquito repellent from *Eucalyptus camaldulensis*. *J Agric Food Chem* 41:2164–2166.

(received 23 July 2015; accepted 1 September 2015)