

# OXYTOCIN INJECTION BEFORE SEMEN COLLECTION INCREASED SEMEN QUANTITY AND QUALITY IN OSTRICHES

Peerasak Suttiyotin<sup>1</sup>, Pacharaporn Khumpim<sup>1\*</sup>, Sakdichod Kimsakulvech<sup>1</sup> and Chalermkwan Eurlaphan<sup>2</sup>

*Received: Jul 18, 2011; Revised: Jan 16, 2012; Accepted: Jan 18, 2012*

## Abstract

The effect of intravenous oxytocin injection before semen collection on semen quality was investigated in ostriches. Six clinically healthy ostriches, aged 3-5 years, previously trained for semen collection by manual stimulation were included in the experiment. These ostriches were routinely used for semen collection. Prior to oxytocin injection, they were subjected to 2 successive semen collections, 4 days apart. Then, they were rested for another 4 days after the second semen collection. On the day of the third semen collection, the birds were intravenously injected with 5 IU of oxytocin via phallic vein. Semen collection was attempted 5 minutes after the injection. Four days after the third semen collection, ejaculates were again obtained from the birds. Seminal characteristics included in the evaluation were semen volume, percentage of motile sperm, sperm concentration, percentages of live, abnormal, and hypo-osmotic swollen spermatozoa.

Semen volume was increased from  $0.27\pm 0.08$  and  $0.23\pm 0.06$  ml (8 and 4 day before injection, respectively) to  $0.95\pm 0.17$  ml ( $p<0.05$ ) on the day of oxytocin injection and returned to  $0.26\pm 0.08$  ml 4 days later. Sperm motility was also increased from  $18.13\pm 9.26$  and  $23.75 \pm 9.76\%$  (8 and 4 day before injection, respectively) to  $57.5\pm 12.54$  ( $p<0.05$ ) on the day of injection and returned to  $23.75\pm 10.68\%$  4 days after injection. Sperm concentration expressed similar results, and higher concentration was obtained on the day of injection ( $753.13\pm 354.94\times 10^6$  cells/ml) compared to 8 and 4 days before and 4 days after injection ( $62.25\pm 31.68$ ,  $143.06\pm 102.91$ , and  $70.78\pm 32.33\times 10^6$  cells/ml, respectively,  $p<0.05$ ). Oxytocin enhanced the percentages of live and membrane integrity in the same manner. The percentage of live spermatozoa was higher with oxytocin injection ( $73.50\pm 12.55\%$ ) than the others (ranged 21.63-44.25%,  $p<0.05$ ). The percentage of swollen sperm (hypo-osmotic swelling test) in the day of oxytocin injection ( $65.25\pm 13.10\%$ ) was higher than the others (ranged 15.13-31.31%,  $p<0.05$ ). No significant effect was found on percentage of abnormal sperm. We concluded that oxytocin injection before semen collection improved most seminal characteristics in ostriches. Whether or not the injection depleted sperm reserve may need further study.

**Keywords:** Oxytocin, semen quality, motility, semen collection, ostrich

---

<sup>1</sup> Faculty of Veterinary Medicine, Mahidol University, Nakorn Prathom 73170, Thailand. Tel.: 0-2441-5242; Fax.: 0-2441-0937; E-mail: neoxstitch@gmail.com

<sup>2</sup> Thai Food Group Co. Ltd., Bangkok, Thailand

\* Corresponding author

## Introduction

Although reproductive system has been intensively investigated in domestic poultry, very few reproductive technologies are applicable for ostriches (*Struthio camelus*). Mating system needs at least 1 male per 2-3 females, the high ratio that does not allow efficient management of reproductive systems to develop for farming. Artificial insemination is one of the major approaches to improve production and enhance reproductive efficiency (Malecki *et al.*, 2008). Semen collection in ostriches has been described and well documented (Soley and Groenewald, 1999; Hemberger, *et al.*, 2001). This allows assisted reproductive technology to be studied and developed both for semen preservation and artificial insemination. Regardless of collection methods, semen quality and quantity are the main purposes to be achieved.

Spermatogenesis in ostriches is more or less similar to that of chicken. There are 2 types of spermatogonia, type A and type B, in seminiferous epithelium of ostriches (Soley and Groenewald, 1999). They have pale nuclei and line on basement membrane. Spermatogonia divide and give rise to primary spermatocytes which enter first meiotic division. They appear in various stages of first meiotic division and are prominent cells in germinal epithelium of seminiferous tubules. Primary spermatocytes divide and give rise to secondary spermatocytes which have relatively large nucleus about 6  $\mu\text{m}$  in diameter. Secondary spermatocytes divide and spermatids are the result of completion of meiotic division. There are 8 stages of spermiogenesis described in ostriches (Soley and Groenewald, 1999). Spermatids change in morphology during these 8 stages and spermatozoa are the final products of the process. The developing germ cells during all stages of spermatogenesis are joined together by intercellular bridges similar to those of birds and mammals.

To maximize spermatozoa harvesting, high frequency of collection may be applied as far as no injury occurs to male ostriches and no dramatic effect occurs to semen quality.

Alternatively, oxytocin has been used to facilitate semen collection, at least in bulls (Palmer *et al.*, 2004), rams (Bozkurt *et al.*, 2007) and men (Rezk *et al.*, 2004). Oxytocin causes contraction of smooth muscle in female although its function in males is unclear. It has important roles, in female mammals, at least in lactation and parturition. In males, it is believed that oxytocin is involved in sexual behavior such as penile erection and copulatory behavior (Carter, 1992; Argiolas and Melis, 2004). Oxytocin induces penile erection via sensitive brain area at paraventricular nucleus of the hypothalamus. In ostriches, anterior pituitary is not directly connected to the hypothalamus as in mammals. Thus, releasing hormones secreted from the base of the hypothalamus is conveyed via blood stream to anterior pituitary gland. The posterior pituitary gland of ostrich, however, is connected to the hypothalamus as in mammals (Soley and Groenewald, 1999). It secretes neurohormones including oxytocin. To the best of our knowledge, very little information concerning use and application of oxytocin for semen collection in ostriches is available. The aim of this study was thus to investigate whether oxytocin injection before semen collection could increase semen quantity and quality.

## Materials and Methods

### Animals

Six months before experiment, ostriches (*Struthio camelus*), aged 3-5 years, have been trained for semen collection by manual stimulation (Figure 1). Six birds were selected from healthy, trained ostriches for this experiment. Those male ostriches were raised under common management system and separated from female ostriches.

Ostriches were routinely used for semen collection. The birds were subjected to 2 successive semen collections, 4 and 8 days apart prior to oxytocin injection. Ostriches rested for another 4 days after the second semen collection. On the day of the third

semen collection, the birds were intravenously injected with 5 IU of oxytocin via phallic vein. To allow full effect of oxytocin Five minutes after the injection, ejaculates were collected from the birds 5 minutes after hormone injection. Four days after the third semen collection, ejaculates were again obtained from the birds. Seminal characteristics included in the evaluation were semen volume, percentage of motile sperm, sperm concentration, percentages of live, and abnormal and hypo-osmotic swollen spermatozoa.

### Semen collection and evaluation

Semen collection following? Soley and Groenewald, 1999; Hemberger *et al.*, 2001) was conducted in the morning in a quiet environment. Each of the ostriches was held in a plucking crush similar to breeding box for artificial insemination in cattle. Soft bag was applied to head of ostriches to calm down. Once the bird was in the device, the collector accessed to the back end. The phallus was retracted, rinsed with warm water and held in position by an assistant. Soft and clean cloth was used for a firmer grip of the phallus. A collector stimulated seminal papillae with fingers to empty semen from ductus deferens. Ejaculation occurred after a few minutes of stimulation and semen ran down the phallic groove where it was collected into a clean glass container.



**Figure 1. Semen collection by manual stimulation in ostriches. The phallus was held in position away from posterior end and a track of white colored semen (arrow) was collected into the beaker**

Immediately after collection, semen was held at room temperature (25°C) before evaluation. Semen sample was subjectively evaluated for sperm progressive motility under light microscope at magnification of 400× (Carl Zeiss Axiostar Plus microscope). To reduce variation, motility evaluation was conducted by the same person in all semen samples and expressed as the percentage of progressively motile sperm cells. The percentages of sperm viability were evaluated after staining with a vital dye, 1.67% eosin Y in 10% nigrosin solution (Chalah and Brillard, 1998; Hancock, 1951). An equal volume of semen sample was mixed with staining solution at 25°C and smear was made after 30 sec of staining. The smear was air dried before evaluation by counting at least 200 cells under a light microscope in the same day of preparation.

Morphology was assessed by staining semen sample with William's stain. Semen sample was smeared on a clean glass slide, air dried and fixed in 95% absolute alcohol for 5 min. The slide was dipped in chloramine T for 1 minute and then dipped a few times in 95% absolute alcohol and in carbolfuchsin stain for 8 min before being washed in running water. The slide was air dried and evaluated for sperm size, shape and characteristics of head and tail under the light microscope at 400×. At least 200 spermatozoa were counted and percentage of abnormal spermatozoa calculated.

Semen concentration was determined by diluting a semen sample with citrate chloramine T solution at the ratio of 1:100. The diluted semen was loaded in counting chambers of a haemocytometer. The sperm cells were counted and concentration calculated. For hypo-osmotic swelling test, 2 µl of semen was mixed with 20 µl of a 100 mOsm solution (containing 0.49 gm of sodium citrate and 0.9 gm of glucose in 100 ml of distilled water). The solution was incubated at 25°C for 1 h before a drop of mixture was placed on a slide, covered with a cover glass and membrane structures of spermatozoa were evaluated under a microscope at 400×.

### Statistical Analysis

Data were tested for normal distribution with Kolmogorov Smirnov one sample test before being analysed. Wilcoxon test was used as non-parametric statistics of data analysis. Differences were considered significant at  $P < 0.05$ . All statistics was performed using SPSS version 17.0 (SPSS Inc., Chicago, IL, USA).

### Results and Discussion

Oxytocin injection improved most seminal characteristics of ostriches and the values returned to base line at day 4 after injection. Semen volume collected by manual stimulation 8 and 4 days (day -8 and -4) before oxytocin injection were  $0.27 \pm 0.08$  and  $0.23 \pm 0.06$  ml respectively ( $p > 0.05$ ; Table 1). The volume increased to  $0.95 \pm 0.17$  ml ( $p < 0.05$ ) on the day of oxytocin injection and returned to  $0.26 \pm 0.08$  ml 4 days after injection. Sperm motility was increased from  $18.13 \pm 9.26$  and  $23.75 \pm 9.76\%$  (day -8 and -4, respectively) to  $57.5 \pm 12.54\%$  ( $p < 0.05$ ) on the day of injection and returned to  $23.75 \pm 10.68\%$  4 days after injection.

Sperm concentration on the day of injection ( $753.13 \pm 354.94 \times 10^6$  cells/ml) was

higher than those of 8 and 4 days before and 4 days after injection ( $62.25 \pm 31.68$ ,  $143.06 \pm 102.91$  and  $70.78 \pm 32.33 \times 10^6$  cells/ml, respectively,  $p < 0.05$ ). The percentage of live spermatozoa was higher with oxytocin injection ( $73.50 \pm 12.55\%$ ) than the others (ranged  $21.63$ – $44.25\%$ ,  $p < 0.05$ ). The percentage of swollen sperm (hypo-osmotic swelling test) in the day of oxytocin injection ( $65.25 \pm 13.10\%$ ) was higher than the others (ranged  $15.13$ – $31.31\%$ ,  $p < 0.05$ ). No significant effect was found on percentage of abnormal sperm.

Techniques for semen collection in emu and ostriches have been developed, such as artificial cloaca with the dummy or the teaser (Malecki, *et al.*, 1997a; Rybnik *et al.*, 2007), and manual stimulation technique (Hemberger *et al.*, 2001; Soley and Groenewald, 1999). The artificial cloaca technique requires dummy or the teaser and training of birds starting at young ages. We used manual stimulation technique because the available ostriches were mature and familiar with human and handling.

There are some approaches to increase sperm production and sperm output in emu and ostriches. Frequency of semen collection has been proved to increase sperm output. In emu, collection at twice daily frequency resulted in higher sperm output than that of once daily (Malecki, *et al.*, 1997b). In ostriches, collection

**Table 1.** Semen volume, % motility, semen concentration and total sperm compared before and after oxytocin administration\*. Data was shown as mean  $\pm$  SE.

	Days before/after oxytocin injection			
	-8	-4	0	4
Volume (ml)	$0.27 \pm 0.08^a$	$0.23 \pm 0.06^a$	$0.95 \pm 0.17^b$	$0.26 \pm 0.08^a$
Motility (%)	$18.13 \pm 9.26^a$	$23.75 \pm 9.76^a$	$57.50 \pm 12.54^b$	$23.75 \pm 10.68^a$
Concentration ( $\times 10^6$ /ml)	$62.25 \pm 31.68^a$	$143.06 \pm 102.91^a$	$753.13 \pm 354.94^b$	$70.78 \pm 32.33^a$
Total sperm ( $\times 10^9$ cells)	$0.01 \pm 0.08^a$	$0.04 \pm 0.03^a$	$0.69 \pm 0.33^b$	$0.02 \pm 0.01^a$
Live sperm (%)	$21.63 \pm 11.07^a$	$31.44 \pm 11.33^a$	$73.50 \pm 12.55^b$	$44.25 \pm 13.14^a$
Hypo-osmotic swelling test (%)	$15.13 \pm 8.43^a$	$25.75 \pm 9.83^a$	$65.25 \pm 13.10^b$	$31.13 \pm 10.41^a$
Abnormal head (%)	$21.65 \pm 10.95$	$9.69 \pm 2.31$	$17.31 \pm 6.64$	$15.75 \pm 6.39$
Abnormal tail (%)	$20.31 \pm 10.16$	$11.44 \pm 3.30$	$13.38 \pm 3.44$	$12.44 \pm 1.24$

\* Different superscripts within the same row demonstrate significant differences ( $p < 0.05$ )

at 6 h interval yielded more sperm output than those of 24 and 48 h intervals, while sperm motility and viability were not affected (Bonato *et al.*, 2011). This approach is safe, useful and applicable when time and facilities are available.

Semen quality in our study was similar to other reports. Semen volume obtained from untreated ostriches (3-5 years) in this study (0.23 to 0.27 ml) was comparable to 4-6 years-old ostriches (0.1 to 1.5 ml) reported by Hemberger *et al.*, (2001) and adult ostriches (0.8 to 1.8 ml; Khumpim *et al.*, 2010). The sperm motility in our study (18.13 to 23.75% in untreated birds) was also comparable to semen of 3-5 years-old ostriches diluted with BPSE (30.42%; Rattanawaree *et al.*, 2010). Sperm concentration of untreated ostriches in our study (62.25 to 143.06×10<sup>6</sup> sperm/ml) was, however, higher than another report (3.9 to 36.0×10<sup>6</sup> sperm/ml; Hemberger *et al.*, 2001). Semen quality of untreated ostriches is thus comparable to those reported by other investigators.

Oxytocin is believed to have both endocrine and paracrine roles in male reproduction. Its roles have been investigated in many species including catfish (Viveiros *et al.*, 2003), bulls (Palmer *et al.*, 2004), rams (Bozkurt *et al.*, 2007) and men (Rezk *et al.*, 2004). We found that oxytocin increased most seminal characteristic in ostriches as well. This is supported by the studies in rabbits (Fjellstrom *et al.*, 1968) and rams (Bozkurt *et al.*, 2007) while there was no such effect in dogs (Traas and Kustritz, 2004). Our results indicate that oxytocin enhances semen quality and quantity in ostriches. It has proved that semen collection in ostrich yields better semen quality and quantity with oxytocin injection prior to collection. Further study may be needed to elucidate mechanism of oxytocin on semen collection in various species.

## Conclusions

It may be concluded oxytocin injection before semen collection in ostriches increases (or improves) most seminal characteristics in

ostriches. This provides an alternative method for efficient harvesting of spermatozoa. Whether or not the injection depleted sperm reserve may need further study.

## Acknowledgements

The authors thank Malai farm for experimental animals and facilities. Research grant was supported by Mahidol University, Thailand.

## References

- Argiolas, A. and Melis, M.R. (2004). The role of oxytocin and the paraventricular nucleus in the sexual behavior of male mammals. *Physiol. Behav.*, 83:309-317.
- Bonato, M., Rybnik, P.K., Malecki, I.A., Cornwallis, C.K., and Cloetea, S.W.P. (2011). Twice daily collection yields greater semen output and does not affect male libido in the ostrich. *Anim. Reprod. Sci.*, 123:258-264.
- Bozkurt, T., Turk, G., and Gur, S. (2007). Effects of exogenous oxytocin on serologic and seminal steroids and semen characteristics in rams. *Turk. J. Vet. Anim. Sci.*, 31:303-309.
- Carter, C.S. (1992). Oxytocin and sexual behavior. *Nerosci. Behav. Rev.*, 16:131-144.
- Chalah, T., and Brillard, J.P. (1998). Comparison of assessment of fowl sperm viability by nigrosin-eosin and dual fluorescence (SYBR-14/PI). *Theriogenology*, 50:487-493.
- Fjellstrom, D., Kihlstrom, J.E. and Melin, P. (1968). The effect of synthetic oxytocin upon seminal characteristics and sexual behaviour in male rabbits. *J. Reprod. Fert.*, 17:207-209.
- Hancock, J.L. (1951). A staining technique for the study of temperature-shock in semen. *Nature*, 167:323-324.
- Hemberger, M.Y., Hospes, R., and Bostedt, H. (2001). Semen collection, examination and spermogram in ostriches. *Reprod. Dom. Anim.*, 36:241-243.
- Khumpim P., Suttiyotin, P. and Kimsakulvech, S. (2010). Semen collection by manual stimulation and semen quality in ostriches. *Proceedings of the International Conference on Veterinary Science*; November 2-5, 2010; Bangkok, Thailand, p. 209-214.
- Malecki, I.A., Martin, G.B., and Lindsay, D.R. (1997a). Semen production by the emu (*Dromaius novaehollandiae*). 1. Methods for collection of semen. *Poultry Sci.*, 76:615-621.
- Malecki, I.A., Martin, G.B., and Lindsay, D.R. (1997b). Semen production by the emu (*Dromaius novaehollandiae*). 2. Effect of

- collection frequency on the production of semen and spermatozoa. *Poultry Sci.*, 76:622-626.
- Malecki, I.A., Rybnik, P.K., and Martin, G.B., (2008). Artificial insemination technology for ratites: a review. *Aust. J. Exp. Agric.*, 48:1284–1292.
- Palmer, C.W., Amundson, S.D., Brito, L.F.C., Waldner, C.L., and Barth, A.D. (2004). Use of oxytocin and cloprostenol to facilitate semen collection by electroejaculation or transrectal massage in bulls. *Anim. Reprod. Sci.*, 80:213-223.
- Rattanawaree, R., Bunyachodmongkol, T., Piboon, L., Suttiyotin, P., Kimsakulvech, S. and Khumpim, P. (2010). Effects of diluents and storage times at 4°C on semen qualities in ostriches. *Proceedings of the International Conference on Veterinary Science*; November 2-5, 2010; Bangkok, Thailand, p. 86-94.
- Rezk, A.Y., Al-Inany, H., and Bedaiwy, M. (2004). Multiple intravenous oxytocin injection may yield sperm in azoospermic men scheduled for TESE. *Middle East Fert. Soc. J.*, 9:238-241.
- Rybnik, P.K., Horbanczuk, J.O., Naranowicz, H., Lukaszewicz, E., and Malecki, I.A. (2007). Semen collection in the ostrich (*Struthio camelus*) using a dummy or a teaser female. *Br. Poultry Sci.*, 48:635–643.
- Soley, J.T., and Groenewald, H.B. (1999). Reproduction. In: *The Ostrich: Biology, Production and Health*. Deeming, D.C. (ed.). CABI Publishing, Oxton, England, p. 129-158.
- Traas, A.M. and Kustritz, M.V. (2004). Effect of administrating oxytocin or prostaglandin F2alpha on characteristics of the canine ejaculate. *Can. Vet. J.*, 45:999-1002.
- Viveiros, A.T.M., Jatzkowski, A., and Komen, J. (2003). Effects of oxtocin on semen release response in African cat fish (*Clarias gariepinus*). *Theriogenology*, 59:1905-1917.