

Subchronic toxicity of *Cissus quadrangularis* Linn.

Aimmanas Attawish¹, Pranee Chavalittumrong², Songpol Chivapat³, Anchalee Chuthaputti⁴, Sadudee Rattanajarasroj⁵ and Somkiat Punyamong⁶

Abstract

Attawish, A., Chavalittumrong, P., Chivapat, S., Chuthaputti, A., Rattanajarasroj, S. and Punyamong, S.

Subchronic toxicity of *Cissus quadrangularis* Linn.

Songklanakarin J. Sci. Technol. 2002, 24(1) : 39-51

Cissus quadrangularis Linn (*C. quadrangularis*), or "Phet-Cha-Sung-Khaat" in Thai, is one of the most commonly used medicinal plants in Thailand for the treatment of hemorrhoid; however, the safety of this herb upon long-term consumption has never been reported. Toxicity study was conducted to evaluate the three-month subchronic toxicity of *C. quadrangularis* powder in five groups of 12 Wistar rats of each sex. Water control group received orally 10 ml of water/kg BW/day. The dried-stems powder was given orally to the four treatment groups at the doses of 0.03, 0.3, 3.0 and 3.0 g/kg BW/day, which were equivalent to 1, 10, 100 and 100 fold of the therapeutic dose in human, respectively, the last group was the recovery group. No difference of initial or final body weights between *C. quadrangularis*-treated and control groups was detected. It was found that *C. quadrangularis* did not produce any significant dose-related changes of hematological parameters or serum clinical chemistry, and no histopathological lesion of any internal organ that could be due to the toxic effect of *C. quadrangularis* was observed. The results indicated that *C. quadrangularis* at the doses given did not produce any toxicity in the rats during the administration period of 3 months.

Key words : *Cissus quadrangularis* Linn., subchronic toxicity

¹M.S. (Nutritional Toxicology), ²M.S. (Phytochemistry), ³M.S. (Pathobiology), ⁴Ph.D. (Pharmacology), ⁵M.S. (Pharmacology), ⁶Cert. in Medical Science Technology, Medicinal Plant Research Institute, Department of Medical Sciences, Ministry of Public Health, Nonthaburi 11000 Thailand.

Corresponding e-mail : aimmanas@dmisc.moph.go.th

Received, 9 October 2000

Accepted, 26 September 2001

บทคัดย่อ

เอมมนัส อัดตวิชัย ปรานี ขวลิตรำรง ทรงพล ชีวะพัฒน์ อัญชลี จูทะพุทธิ สดุดี รัตนจรัสโรจน์
และ สมเกียรติ ปัญญามัง

พิษกึ่งเรื้อรังของเพชรสังฆาต (*Cissus quadrangularis* Linn.)

ว. สงขลานครินทร์ วทท. 2545 24(1) : 39-51

เพชรสังฆาต (*Cissus quadrangularis* Linn.) เป็นสมุนไพรไทยที่ใช้กันมากอีกชนิดหนึ่ง ในการบำบัดอาการริดสีดวงทวาร โดยที่ไม่มียารายงานความปลอดภัยของสมุนไพรนี้หากใช้เป็นเวลานาน ดังนั้นคณะผู้วิจัยจึงได้ศึกษาพิษกึ่งเรื้อรัง (3 เดือน) ของเพชรสังฆาต ในหนูขาวพันธุ์วีสตาร์ 5 กลุ่ม ๆ ละ 12 ตัว/เพศ กลุ่มควบคุมได้รับน้ำทางปาก 10 มล./น้ำหนักตัว 1 กก./วัน ขณะที่หนูอีก 4 กลุ่มได้รับผงยาเพชรสังฆาตแห้งทางปากในขนาด 0.03, 0.3, 3.0 และ 30 กรัม/น้ำหนักตัว 1 กก./วัน หรือเทียบเท่า 1, 10, 100 และ 100 เท่าของขนาดที่ใช้ในคน/วัน ตามลำดับ โดยกลุ่มสุดท้ายเป็นกลุ่มสังเกตอาการหลังการหยุดยา ผลการศึกษาพบว่า การเจริญเติบโตของกลุ่มที่ได้รับผงยาและกลุ่มควบคุมไม่แตกต่างกัน ไม่ทำให้เกิดการเปลี่ยนแปลงของค่าทางโลหิตวิทยาและค่าทางชีวเคมี หรือจุดพยาธิสภาพของอวัยวะภายในที่มีความสัมพันธ์กับขนาดของผงยา และไม่พบความผิดปกติใด ๆ ที่สามารถสรุปได้ว่าเนื่องมาจากความเป็นพิษของผงยาเพชรสังฆาต จึงอาจกล่าวได้ว่าเพชรสังฆาตในขนาดและระยะเวลาดังกล่าวไม่มีพิษต่อหนูขาว

สถาบันวิจัยสมุนไพร กรมวิทยาศาสตร์การแพทย์ กระทรวงสาธารณสุข อำเภอเมือง จังหวัดนนทบุรี 11000

The use of medicinal plants is an alternative way for Thai people to take care of their own health because of their low cost, easy access and, most of all, their use since ancient times. *Cissus quadrangularis* Linn. (*C. quadrangularis*), commonly known in Thailand as "Phet-Cha-Sung-Khaat", is a climber that belongs to the Vitaceae family. It is one among the most frequently used medicinal plants in Thailand which can be found throughout the country (Pongboonrod, 1950). The stout fleshy quadrangular stem of *C. quadrangularis* is traditionally used for treatment of hemorrhoid and irregular menstruation and also used for acceleration of bone fracture healing. (Banjob, et al., 2000a; Tiangburanatham, 1996; Wuthithammawej, 1994; Quisumbing, 1951; Banjob, et al., 2000b; Chopra, et al., 1976; Chopra, et al., 1975; Udupa and Prasad, 1964). Tiangburanatham (1996) has reported that 100 g of fresh stem contained about 267 mg of carotene, 398 mg of vitamin C and a lot of calcium oxalate crystals. Phytochemical constituents contained triterpene, i.e., α -amyrin and α -amyrone (Bhutani, et al., 1984), phytosterol, i.e., β -sitosterol (Pluemjai and Saifah, 1986), steroid,

i.e., keto-steroid (Udupa, et al., 1965), oxo-steroid (Sen, 1964) and two new unsymmetric tetracyclic triterpenoids, namely, onocer-7-ene-3 α , 21 β -diol and onocer-7-ene-3 β , 21 α -diol (Madan and Ram, 1990). Furthermore, it was recently found that three new stilbene derivatives, quadrangularins A, B, and C, could be isolated from the stems of *C. quadrangularis* together with resveratrol, piceatannol, pallidol and parthenocissine A (Saburi, et al., 1999).

The project of 'Traditional medicines and primary health care' was undertaken by the Ministry of Public Health to bring about the scientific proof of *C. quadrangularis* as a hemorrhoid remedy. Accordingly, Lhieochaiphunt and Sangdee (1985) have studied the acute toxicity of aqueous crude extract in mice, rats and rabbits. The results showed that it was slightly toxic to practically nontoxic depending on the route of administration. Segsunviriya and Choomprabutra (1989) conducted a clinical trial of *C. quadrangularis* for the treatment of hemorrhoid and found that it was effective. A recent report of acute toxicity in ICR mice demonstrated that it was non toxic. In subchronic

toxicity testing (30 days) in Wistar rats, it was found that *C. quadrangularis* caused low levels of serum creatinine, albumin and WBC. These occurrences disappeared within 14 days when the rats were allowed to recover (Limpanussorn, *et al.*, 2000). In addition, the Chao-Praya-Apaipubate General Hospital in Prachinburi province, the leading hospital in utilizing herbal medicines for the treatment of common diseases, is dispensing the dried stem powder in the dose of 1.5-3.0 g for the treatment of hemorrhoid. However, no report for long-term toxicity test has been made. Thus in order to evaluate the safety of this plant, the present study, using an OECD guideline (1998), was conducted to investigate the subchronic toxicity of *C. quadrangularis* when administered orally for three months in rats.

Materials and Methods

Plant material and preparation

The plant was collected from the botanical garden of Chao-Praya-Apaipubate Hospital in Prachinburi province, and was verified by Miss Supaporn Pitiporn, the head pharmacist of the hospital. A voucher specimen (BKF 092329) was deposited at the Forest Herbarium, Royal Forest Department, Bangkok, Thailand.

The plant was washed with tap water, cut, dried in a hot air oven at 50°C, ground and sieved with sieve No.100, then the powder stored in a brown bottle with cap at room temperature. The powder was suspended to the desired concentrations with distilled water. Phytochemical determination showed the quercetin constituent equivalent to 0.1216 microgram percent.

Treatment of the animals

Sixty male Wistar rats weighing 210 ± 10 g and 60 female rats weighing 170 ± 10 g from the National Laboratory Animal Center, Mahidol University, Nakornpathom province, were used. The animals were housed in the animal facility of the Department of Medical Sciences. The temperature in the animal room was kept at $25 \pm 1^\circ\text{C}$ with 60% relative humidity. The animals were allowed

to have free access to food and clean water.

Three months toxicity study

According to the OECD guideline (1998), sixty Wistar rats of each sex were randomly divided into 5 groups of 12 animals per sex. Group 1 (water control) received water 10 milliliter/kilogram of body weight/day (ml/kg BW/day). Groups 2-5 were given the suspensions which were equivalent to 0.03, 0.3, 3.0 and 3.0 grams of dried powdered plant /kilogram of body weight/day (g/kg BW/day), which were equivalent to 1, 10, 100 and 100 folds of therapeutic dose (1.5 g/50-kg person/day), respectively. The last group was the recovery group. Body weight and food-intake were measured weekly and the animals were observed for signs of abnormalities for 3 months. At the end of the treatment period, the 1st - 4th groups of rats were fasted for 18 hours, then anesthetized with ether and sacrificed by drawing blood samples from the inferior vena cava for hematological and biochemical examinations. The 5th group of rats was withdrawn from of the feeding of the plant for another 14 days before being sacrificed.

Hematological analysis was performed using an automatic hematological analyzer (Cell Dyne 3500, Abbott). The parameters of the blood samples measured were: hematocrit (Hct), hemoglobin (Hb), red blood cell (RBC), mean cell volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC), red cell distribution width (RDW), white blood cell (WBC), % neutrophil (%N), % lymphocyte (%L), % monocyte (%M), % eosinophil (%E), % basophil (%B), platelet, mean platelet volume (MPV), plateletcrit (PCT), platelet distribution width (PDW) and reticulocyte.

Biochemical analysis of serum samples was performed using an automatic chemistry analyzer (Hitachi model 912). Biochemical parameters measured were alkaline phosphatase (ALP), alanine aminotransferase (ALT), aspartate aminotransferase (AST), blood urea nitrogen (BUN), p-amylase, total protein, albumin, bilirubin, creatinine, glucose, uric acid, triglyceride, cholesterol, sodium, potassium and chloride.

The positions, shapes, sizes and colors of internal organs, namely, brain, heart, both kidneys and lungs, trachea, esophagus, stomach, liver, pancreas, intestine, spleen, bladder, salivary gland, adrenal gland and testis in male rats or ovary and uterus in female rats were visually observed for any signs of gross lesions. These organs were then collected, weighed to determine relative organ weights, and preserved in 10% phosphate buffered formalin solution. Tissue slides were prepared and stained with hematoxylin and eosin and histopathological examinations were performed by a veterinary pathologist.

Statistical Analysis

The data were analyzed by one-way ANOVA followed by Duncan multiple range test, using SPSS/PC program, to determine significant differences between groups at $p < 0.05$. Histopathological data were evaluated by the Fisher exact test and the significance level was set at $p < 0.05$.

Results and Discussion

Effects of the *C. quadrangularis* on body weight, food intake and relative organ weight

In both male and female animals, there was no difference in the average body weights between *C. quadrangularis*-treated groups and control group throughout the experimental period of 3 months (Figure 1). It was found that food consumption of animals receiving the *C. quadrangularis* was significantly different from the control groups for several weeks. Male rats receiving *C. quadrangularis* 3.0 g/kg/day had lower food intake than the male control group during 4th - 12th weeks and female rats receiving *C. quadrangularis* 0.03 g/kg/day had higher food intake than the female control group during 1st - 13th weeks of the study (Figure 2). In male rats, there was no difference in the relative organ weight between *C. quadrangularis*-treated group and control group (Table 1). Female rats treated with *C. quadrangularis* at the dose of 0.03 g/kg/day had lower relative organ weights of the brain and heart than the control group (Table 2).




Figure 1. Growth curves of male and female rats receiving *C. quadrangularis* for 3 months




Figure 2. Food consumption of male and female rats receiving *C. quadrangularis* for 3 months

Table 1. Body weight (g) and relative organ weight (g/kg) of male rats receiving *C. quadrangularis* for 3 months

Organs	Dose of <i>C. quadrangularis</i> (g/kg BW/day)				
	control	0.03	0.3	3.0	3.0-R
	n = 12	n = 12	n = 12	n = 12	n = 12
Initial body weight	205±12	211±10	208±10	208±9	202±7
Final body weight	493±37	502±47	498±29	485±37	482±34
Brain	4.28±0.32	4.21±0.36	4.27±0.30	4.26±0.18	4.25±0.36
Heart	2.62±0.18	2.55±0.25	2.69±0.20	2.55±0.18	2.60±0.30
Lung	3.57±0.38	3.51±0.32	3.38±0.29	3.52±0.32	3.29±0.28
Liver	25.54±1.19	25.56±1.09	25.55±2.48	25.75±1.78	25.18±1.70
Stomach	3.72±0.32	3.73±0.35	3.77±0.39	3.88±0.37	3.65±0.20
Spleen	1.87±0.20	1.88±0.18	1.88±0.18	1.89±0.14	1.88±0.26
Right Kidney	2.58±0.18	2.55±0.14	2.64±0.17	2.61±0.18	2.59±0.19
Left Kidney	2.51±0.18	2.47±0.16	2.52±0.15	2.50±0.17	2.48±0.16
Right Testis	6.61±0.42	6.52±0.57	6.36±0.41	6.78±0.69	6.45±0.70
Left Testis	6.53±0.41	6.46±0.45	6.26±0.34	6.59±0.86	6.30±0.65
Right Adrenal	0.079±0.012	0.077±0.013	0.080±0.015	0.076±0.014	0.077±0.013
Left Adrenal	0.083±0.014	0.082±0.018	0.084±0.016	0.081±0.013	0.082±0.024
Bladder	0.255±0.042	0.274±0.035	0.240±0.042	0.262±0.037	0.238±0.054

R: Recovery group

The values are expressed as mean ± SD.

Table 2. Body weight (g) and relative organ weight (g/kg) of female rats receiving *C. quadrangularis* for 3 months

Organs	Dose of <i>C. quadrangularis</i> (g/kg BW/day)				
	control	0.03	0.3	3.0	3.0-R
	n = 12	n = 12	n = 12	n = 12	n = 12
Initial body weight	176±9	175±10	176±7	171±6	172±9
Final body weight	292±20	306±26	294±18	284±22	291±23
Brain	6.82±0.45	6.34±0.51*	6.71±0.44	6.91±0.54	6.75±0.62
Heart	3.06±0.27	2.81±0.22*	3.08±0.27	2.87±0.25	3.09±0.39
Lung	4.46±0.42	4.54±0.22	4.28±0.30	4.57±0.47	4.23±0.42
Liver	23.84±2.98	24.57±1.82	24.19±1.95	24.57±1.71	24.23±3.32
Stomach	4.98±0.76	4.89±0.60	4.94±0.32	5.14±0.50	5.11±0.59
Spleen	2.37±0.27	2.20±0.22	2.43±0.24	2.41±0.33	2.49±0.31
Right Kidney	2.84±0.24	2.78±0.20	2.79±0.27	2.78±0.25	2.86±0.32
Left Kidney	2.71±0.23	2.59±0.21	2.69±0.27	2.66±0.23	2.66±0.25
Right Adrenal	0.148±0.037	0.146±0.028	0.160±0.032	0.144±0.019	0.138±0.024
Left Adrenal	0.163±0.042	0.148±0.033	0.165±0.027	0.149±0.026	0.153±0.024
Uterus	3.00±0.75	2.80±1.06	3.03±0.56	2.50±0.31	2.96±0.75
Bladder	0.272±0.058	0.295±0.060	0.274±0.064	0.280±0.059	0.246±0.029

* significantly different from control group (p<0.05)

R: Recovery group

The values are expressed as mean ± SD.

Effect of *C. quadrangularis* on hematological parameters

There was no difference of hematocrit, the number of red blood cells, hemoglobin, MCV, MCH, MCHC, the number of white blood cell, %N, %L, platelet, PCT, PDW, MPV or the number of reticulocyte between *C. quadrangularis*-treated groups and control groups of either male or female rats (Tables 3 and 4). However the groups of male rats receiving *C. quadrangularis* had significantly lower %B than that of the control. Male rats receiving *C. quadrangularis* at the dose of 3.0 g/kg/day had significantly lower %M than that of the control. Male rats receiving *C. quadrangularis* at the dose of 0.3 g/kg/day had significantly lower percent of reticulocyte than that of the control but these values were within normal range (Gad, 1992). Female rats receiving *C. quadrangularis* at the dose of 3.0 g/kg/day had significantly higher

%E than that of the control but these values were within normal range (Gad, 1992). In recovery groups, it was found that *C. quadrangularis* caused high %N and low %B in male, but caused high %B and low MCHC in female, however these values were within normal range (Gad, 1992).

Effect of the *C. quadrangularis* on blood chemistry

In male and female rats, no difference in the serum levels of ALP, ALT, AST, BUN, p-amylase, total protein, albumin, bilirubin, creatinine, glucose, uric acid, triglyceride, cholesterol, sodium, potassium and chloride was found between *C. quadrangularis*-treated groups and the control groups. It was found that *C. quadrangularis* caused high level of serum bilirubin, but caused low level of serum uric acid in male rats which were allowed to recover (Tables 5 and 6), however the values were

Table 3. Hematological values of male rats receiving *C. quadrangularis* for 3 months

Parameters	Dose of <i>C. quadrangularis</i> (g/kg BW/day)				
	control	0.03	0.3	3.0	3.0-R
	n = 12	n = 12	n = 12	n = 12	n = 12
Hematocrit (%)	45.70±3.04	45.64±2.22	46.30±1.36	44.95±3.33	44.99±2.37
RBC (x10⁶ cells/mm³)	9.11±0.71	8.90±0.34	9.11±0.34	8.84±0.39	8.78±0.45
Hemoglobin (g/dl)	15.68±0.55	15.62±0.59	15.62±0.40	15.48±0.48	15.45±0.52
MCV (µm³/red cell)	50.18±1.81	51.25±1.76	50.85±1.55	50.79±2.40	51.25±1.03
MCH (pg/red cell)	17.30±1.22	17.55±0.54	17.15±0.39	17.57±0.62	17.63±0.69
MCHC (g/dl RBC)	34.52±2.51	34.28±1.13	33.75±0.56	34.72±2.29	34.43±1.53
WBC (x10³ cells/mm³)	5.62±1.38	5.70±1.28	5.90±1.26	5.86±0.91	5.18±0.79
Neutrophil (%)	11.72±2.89	11.17±3.01	13.58±4.3	13.34±3.83	15.53±6.35*
Eosinophil (%)	1.60±0.73	1.37±0.63	1.78±0.66	2.07±0.64	1.74±0.67
Lymphocyte (%)	81.77±5.55	84.25±4.36	81.07±5.23	82.17±3.65	79.07±6.66
Monocyte (%)	3.09±2.42	1.94±1.49	2.39±1.27	1.33±1.43*	2.44±1.26
Basophil (%)	1.83±1.19	1.27±0.30*	1.17±0.27*	1.10±0.44*	1.21±0.39*
Platelet (x10³ cells/mm³)	949±140	866±67	909±100	913±93	865±77
PCT (%)	0.86±0.20	0.79±0.16	0.86±0.11	0.76±0.17	0.86±0.11
PDW (%CV)	17.57±2.61	17.51±2.79	18.29±0.41	16.60±3.62	18.56±0.66
MPV (fl/platelet)	9.06±1.47	9.04±1.54	9.42±0.51	8.53±1.94	9.93±1.15
Reticulocyte (x10³ cells/mm³)	235±43	234±41	205±29	204±33	245±30
Reticulocyte (%)	2.61±0.39	2.62±0.46	2.26±0.33*	2.34±0.36	2.81±0.25

* significantly different from control group (p<0.05)

R: Recovery group

The values are expressed as mean ± SD.

within normal range (Gad, 1992). The significant change of serum sodium in recovery groups was less than that of the control group, however the lowered serum sodium had no clinical effect. This

change did not occur in any of the treated groups. Hence, this change may not be due to the effect of *C. quadrangularis*.

Table 4. Hematological values of female rats receiving *C. quadrangularis* for 3 months

Parameters	Dose of <i>C. quadrangularis</i> (g/kg BW/day)				
	control	0.03	0.3	3.0	3.0-R
	n = 12	n = 12	n = 12	n = 12	n = 12
Hematocrit (%)	44.52±2.35	44.12±1.77	45.02±1.65	44.30±1.98	45.54±2.28
RBC (x10 ⁶ cells/mm ³)	8.02±0.41	7.88±0.30	8.13±0.30	8.08±0.42	8.13±0.22
Hemoglobin (g/dl)	15.01±0.51	14.78±0.71	15.16±0.51	14.86±0.42	14.98±0.76
MCV (xm ³ /red cell)	55.54±1.93	56.03±2.07	55.46±2.96	54.91±1.39	56.02±1.97
MCH (pg/red cell)	18.75±0.73	18.78±0.91	18.70±1.05	18.48±0.63	18.44±0.76
MCHC (g/dl RBC)	33.77±0.89	33.51±0.79	33.70±0.52	33.66±0.82	32.91±0.50*
WBC (x10 ³ cells/mm ³)	2.66±0.65	2.79±0.53	2.89±0.87	2.71±0.57	2.40±0.47
Neutrophil (%)	10.99±4.00	14.77±5.22	12.20±4.59	14.61±6.02	14.85±3.65
Eosinophil (%)	1.60±0.31	1.53±0.47	1.88±0.85	2.27±1.10*	1.53±0.44
Lymphocyte (%)	84.53±4.12	80.12±6.83	83.27±5.86	79.62±6.79	79.22±5.37
Monocyte (%)	1.93±0.89	2.77±2.07	1.78±1.80	2.58±1.61	3.10±2.16
Basophil (%)	0.96±0.43	0.82±0.31	0.86±0.41	0.94±0.36	1.30±0.48*
Platelet (x10 ³ cells/mm ³)	868±54	872±88	919±103	873±61	883±75
PCT (%)	0.79±0.05	0.80±0.08	0.85±0.08	0.80±0.07	0.83±0.06
PDW (%CV)	18.01±0.46	18.20±0.22	18.15±0.43	18.07±0.31	18.00±0.21
MPV (fl/platelet)	9.18±0.59	9.20±0.49	9.25±0.60	9.11±0.43	9.43±0.38
Reticulocyte (x10 ³ cells/mm ³)	261±65	275±59	264±75	240±45	298±70
Reticulocyte (%)	3.29±0.79	3.46±0.72	3.24±0.88	2.98±0.53	3.66±0.83

* significantly different from control group (p<0.05)

R: Recovery group

The values are expressed as mean ± SD.

Effect of the *C. quadrangularis* on histopathology of internal organs

Upon gross examination of internal organs, no abnormal signs were observed. Histopathologi-

cal results indicated that there was no lesion of salivary gland, spleen, pancreas or intestine in any group of animals (Tables 7 and 8). The incidence of peribronchiolar lymphoid aggregation in all

Table 5. Biochemical values of male rats receiving *C. quadrangularis* for 3 months

Parameters	Dose of <i>C. quadrangularis</i> (g/kg BW/day)				
	control	0.03	0.3	3.0	3.0-R
	n = 12	n = 12	n = 12	n = 12	n = 12
ALP (U/L)	64.00±11.39	64.08±8.94	65.92±8.73	60.17±8.17	66.50±8.02
ALT (U/L)	32.83±8.32	31.67±5.42	34.75±7.42	38.50±6.29	38.67±14.64
AST (U/L)	71.42±11.44	8.33±4.72	65.08±8.34	65.92±10.09	66.25±10.70
p-amylase (U/L)	1989±266	1960±185	2154±288	2070±205	1958±216
Total protein (g/dl)	6.57±0.76	6.50±0.32	6.66±0.31	6.49±0.42	6.72±0.22
Albumin (g/dl)	3.33±0.40	3.35±0.23	3.38±0.23	3.32±0.24	3.23±0.08
Bilirubin (mg/dl)	0.07±0.03	0.07±0.02	0.07±0.02	0.06±0.03	0.11±0.03*
BUN (mg/dl)	18.32±2.55	17.18±2.29	19.49±3.41	18.60±2.97	19.38±3.19
Creatinine (mg/dl)	0.64±0.07	0.64±0.04	0.65±0.05	0.62±0.06	0.62±0.04
Glucose (mg/dl)	170.48±25.77	169.16±18.26	175.09±23.06	170.54±18.18	157.54±11.48
Uric acid (mg/dl)	2.12±1.05	1.64±0.58	1.94±0.62	1.80±0.71	1.42±0.43*
Triglyceride (mg/dl)	127.54±46.73	125.72±38.37	144.11±33.39	145.38±48.91	131.76±12.78
Cholesterol (mg/dl)	72.39±24.92	62.24±8.62	70.04±10.98	67.13±15.03	59.36±12.78
Na ⁺ (mmol/l)	148±3	144±8	146±5	145±6	140±1*
K ⁺ (mmol/l)	5.72±1.12	5.22±0.53	5.30±0.55	5.32±0.51	5.88±0.62
Cl ⁻ (mmol/l)	108±4	106±5	106±2	107±3	110±2

* significantly different from control group (p<0.05)

R: Recovery group

The values are expressed as mean ± SD.

treated groups was significantly lower than the control group, and this incidence was increased in the recovery group. Hence, this change should be due to the effect of *C. quadrangularis*. Park and his colleague (2001) reported that β -sitosterol from the ethanol extract of cactus (*Opuntia ficus indica*) showed anti-inflammatory action in adjuvant-induced chronic inflammation model in mice. Because of β -sitosterol the same constituent in *C. quadrangularis*, this plant may have anti-inflammatory action too. In male rats receiving *C.*

quadrangularis at the dose of 0.03 g/kg/day, the incidence of hepatocyte degeneration of the liver was not significantly different from that of the control. Cortex fatty degeneration of adrenal gland was noted in half of the male control group. However the incidence of fatty degeneration of the adrenal gland in all groups of male rats receiving *C. quadrangularis* was significantly lower than that of control, and there was no difference of the incidence between recovery and control group. This incidence seemed to be sex-related (Gad, 1992).

Table 6. Biochemical values of female rats receiving *C. quadrangularis* for 3 months

Parameters	Dose of <i>C. quadrangularis</i> (g/kg BW/day)				
	control	0.03	0.3	3.0	3.0-R
	n = 12	n = 12	n = 12	n = 12	n = 12
ALP (U/L)	28.42±3.87	28.17±5.37	29.17±5.49	27.25±4.35	27.92±6.57
ALT (U/L)	26.50±5.18	28.75±6.18	26.42±5.68	29.08±5.87	27.83±5.69
AST (U/L)	64.08±8.35	67.00±9.12	63.08±6.96	64.17±10.61	61.58±6.23
p-amylase (U/L)	1150±242	1168±180	1126±200	1081±183	1054±284
Total protein (g/dl)	6.92±0.40	6.91±0.36	6.72±0.42	6.61±0.58	6.98±0.25
Albumin (g/dl)	3.71±0.17	3.69±0.26	3.65±0.26	3.61±0.42	3.54±0.14
Bilirubin (mg/dl)	0.10±0.04	0.10±0.06	0.10±0.04	0.09±0.07	0.09±0.03
BUN (mg/dl)	19.48±1.88	19.74±2.17	19.56±3.04	20.19±3.55	19.43±3.70
Creatinine (mg/dl)	0.71±0.05	0.72±0.04	0.67±0.06	0.70±0.09	0.69±0.04
Glucose (mg/dl)	145.16±14.78	147.90±16.33	128.92±21.28	132.21±28.77	132.82±17.49
Uric acid (mg/dl)	1.62±0.50	1.37±0.50	1.37±0.50	1.48±0.57	1.42±0.61
Triglyceride (mg/dl)	79.05±32.62	73.85±23.84	79.58±38.35	86.26±26.42	84.15±32.48
Cholesterol (mg/dl)	71.71±22.00	73.85±17.98	68.07±12.91	67.99±12.77	67.10±14.15
Na ⁺ (mmol/l)	150±3	148±4	146±6	146±10	141±1*
K ⁺ (mmol/l)	5.00±0.66	4.79±0.60	4.88±1.01	4.61±0.90	4.99±0.68
Cl ⁻ (mmol/l)	114±4	112±4	111±2	110±8	113±2

* significantly different from control group (p<0.05)

R: Recovery group

The values are expressed as mean ± SD.

Therefore, *C. quadrangularis* may reduce fat accumulation in the adrenal cortex. Fraser (1994) reported that β -sitosterol inhibited the absorption of both endogenous and exogenous cholesterol and in moderate doses lowered serum cholesterol. As in female rats receiving *C. quadrangularis*, though there were some lesions detected microscopically in the liver and kidneys in some groups of animals, but this change was not significantly different from control group. We found that only the congestion of adrenal gland in recovery group was signifi-

cantly higher than that of control; however, this lesion might be caused by the stress from anesthetic process as well (Koplewitz, *et al.*, 1998; Pugachev, 1977; Bassett and Cairncross, 1975).

Conclusion

Three-month subchronic toxicity study of *C. quadrangularis* Linn. in Wistar rats indicated that the dried stem powder at the doses of 0.03, 0.3 and 3.0 g/kg BW/day, which were equivalent to 1, 10

Table 7. Histopathological values of male rats receiving *C. quadrangularis* for 3 months

Organs	Microscopic findings	Dose of <i>C. quadrangularis</i> (g/kg BW/day)				
		control	0.03	0.3	3.0	3.0-R
		n = 12	n = 12	n = 12	n = 12	n = 12
Lung	Peribronchiolar lymphoid aggregation	7/12	0/12*	0/12*	2/12*	3/12
Heart	Focal myocarditis	1/12	0/12	0/12	0/12	1/12
Liver	Hepatocyte degeneration	1/12	1/12	0/12	0/12	0/12
Kidney		0/12	0/12	0/12	0/12	0/12
Spleen		0/12	0/12	0/12	0/12	0/12
Pancreas		0/12	0/12	0/12	0/12	0/12
GI tract		0/12	0/12	0/12	0/12	0/12
Testis		0/12	0/12	0/12	0/12	0/12
Adrenal gland	Cortex fatty degeneration	6/12	0/12*	0/12*	1/12*	6/12
Salivary gland		0/12	0/12	0/12	0/12	0/12

* significantly different from control group (p<0.05)

R: Recovery group

The results are expressed as number of rats with pathological findings/ total number of rats examined.

and 100 fold the therapeutic dose did not produce any significant dose-related changes of hematological parameters, serum biochemistry or histopathology of any internal organs. Therefore, it is concluded that *C. quadrangularis* at the given doses did not produce any significant toxic effect in rats during the period of treatment for 3 months.

Acknowledgments

The authors are grateful to Dr. Anuthep Rangripipat, Department of Veterinary Pathology, Faculty of Veterinary Science, Chulalongkorn University, for histopathological examinations of tissue samples. Thanks are also due to Miss Supaporn Pittiporn, the head pharmacist of Chao-Praya-Apaipubate Hospital, Prachin Buri province, who identified and supplied dried stems of *C. quadrangularis*, and the staff of the Animal Facility of the

Department of Medical Sciences for the animal care.

References

- Banjob, M., Pecharaply, D., Chaiyaraj, S. and Sawangwong, C. 2000a. Medicinal plants in North Eastern, Part 2. Institute of Medicinal Plant Research, Department of Medical Sciences, Ministry of Public Health. Thanasinjaroen Printing, Bangkok, Thailand. p.55. (in Thai)
- Banjob, M., Pecharaply, D., Chaiyaraj, S. and Sawangwong, C. 2000b. Medicinal plants in North Eastern, Part 3. Institute of Medicinal Plant Research, Department of Medical Sciences, Ministry of Public Health. S.R. Printing Massproducts Co. Ltd., Bangkok, Thailand. p.68. (in Thai)
- Bassett, J.R. and Cairncross, K.D. 1975. Morphological changes induced in rats following prolonged exposure to stress. *Pharmacol. Biochem. Behav.* 3(3): 411-20.

Table 8. Histopathological values of female rats receiving *C. quadrangularis* for 3 months

Organs	Microscopic findings	Dose of <i>C. quadrangularis</i> (g/kg BW/day)				
		control	0.03	0.3	3.0	3.0-R
		n = 12	n = 12	n = 12	n = 12	n = 12
Lung	Peribronchiolar lymphoid proliferation	3/12	0/12*	0/12*	1/12*	2/12
Heart	Focal myocarditis	0/12	0/12	0/12	0/12	0/12
Liver	Lymphoid aggregated periportal area	0/12	1/12	1/12	0/12	1/12 (focal necrosis)
Kidney	Tubular cast cyst	7/12	9/12	4/12	6/12	7/12
	Lymphoid aggregation	0/12	0/12	0/12	0/12	0/12
Spleen		0/12	0/12	0/12	0/12	0/12
Pancreas		0/12	0/12	0/12	0/12	0/12
GI tract		0/12	0/12	0/12	0/12	0/12
Ovary		0/12	0/12	0/12	0/12	0/12
Uterus						
Cervix						
Adrenal gland	Congestion	0/12	0/12	0/12	0/12	6/12*
Salivary gland		0/12	0/12	0/12	0/12	0/12

* significantly different from control group ($p < 0.05$)

R: Recovery group

The results are expressed as number of rats with pathological findings/total number of rats examined.

- Bhutani, K.K., Kapoor, R. and Atal, C.K. 1984. Two unsymmetric tetracyclic triterpenoids from *Cissus quadrangularis*. *Phytochemistry*. 23(2): 407-10.
- Chopra, S.S., Patel, M.R. and Awadhiya, R.P. 1976. Studies on *Cissus quadrangularis* in experimental fracture repair: a histopathological study. *Indian J. Med. Res.* 64:1365-75.
- Chopra, S.S., Patel, M.R., Gupta, L.P. and Datta, I.C. 1975. Studies on *Cissus quadrangularis* in experimental fracture repair: effect on chemical parameters in blood. *Indian J. Med. Res.* 63:824-8.

- Fraser, GE. 1994. Diet and coronary heart disease: beyond dietary fats and low-density-lipoprotein cholesterol. *Amer. J. Clin. Nutri.* 59: 1117-23.
- Gad, S.C. 1992. The Rat: Pathology. In: Gad, S.C., Chengellis, C.P. (Eds.), *Animal Model in Toxicology*, Marcel Dekker, New York, USA. pp. 81, 111-3.
- Koplewitz, B.Z., Daneman, A., Cutz, E. and Hellmann, J. 1998. Neonatal adrenal congestion: a sonographic-pathologic correlation. *Pediatr. Radiol.* 28(12): 958-62.
- Lhieochaiphunt, S. and Sangdee, P. 1985. An acute toxicity testing of *Cissus quadrangularis*;

- Vitaceae. Research Report. Faculty of Pharmacy, Chiang Mai University. pp.15. (in Thai)
- Limpanussorn, J., Poongchompu, S., Thisayakorn, K. *et al.* 2000. Toxicity study of *Cissus quadrangularis* L. In: Guideline for Medicinal Plant Development of Thailand. Thailand Institute of Scientific and Technological Research. p.47. (in Thai)
- Madan, M.G. and Ram, K.V. 1990. Unsymmetric tetracyclic triterpenoid from *Cissus quadrangularis*. *Phytochemistry*. 29(1): 336-7.
- OECD. 1998. OECD Principles of Good Laboratory Practice, Section 2. OECD, Paris, France.
- Park, E., Kahng, J., Lee, S.H., Shin, K. 2001. An anti-inflammatory principle from cactus. *Fitoterapia*. 72(3): 288-90.
- Pongboonrod, S. 1950. Mai Thed Muang Thai. Kaseambunnakit Printing, Bangkok, Thailand. pp.428-9. (in Thai)
- Pugachev, M.K. 1977. Problem of thickening of the adrenal cortex in acute thermal stress. *Arkh. Anat. Gistol. Embriol.* 72(3): 73-7.
- Quisumbing, E. 1951. Medicinal plants of the Philippines. *Tech. Bull.* 16: 1-10.
- Saburi, A.A., Rene, N., Marie-Therese, M., *et al.* 1999. Stilbene derivatives from *Cissus quadrangularis*. *J. Nat Prod.* 62: 1694-5.
- Segsunviriya, C. and Choomprabutra, S. 1989. A clinical study of *Cissus quadrangularis* Linn. in hemorrhoid patients. Seminar of Research and Development of Medicinal Plant. Division of Medical Research, Department of Medical Sciences, Ministry of Public Health. pp. 54-5. (in Thai)
- Sen, S.P. 1964. Active constituents (oxo steroids) of *Cissus quadrangularis*. *Indian J. Pharmacy*. 26(9): 247-8.
- Tiangburanatham, W. 1996. Dictionary of Thai Medicinal Plants. Prachumtong Printing, Bangkok, Thailand. pp. 572-3. (in Thai)
- Pluemjai, T. and Saifah, E. 1986. Constituents of *Cissus quadrangularis* Linn. *Th. J. Pharm.Sci.* 11(4): 205-11.
- Udupa, K.N. and Prasad, G. 1964. Biomechanical and 45-CA studies on the effect of *Cissus quadrangularis* in fracture repair. *Indian J. Med. Res.* 52(5): 480-7.
- Udupa, K.N., Prasad, G and Sen, S.P. 1965. The effect of phytogenic anabolic steroid in the acceleration of fracture repair. *Life Sci.* 4(3): 317-27.
- Wuthithammawej, W. 1994. Thai Traditional Medicine. O.S. Printing House, Bangkok, Thailand. p. 337. (in Thai)