

ไคติเนสและการยับยั้งราก่อโรคจากแอคติโนมัยซีทเอนโดไฟต์

Chitinolytic and antifungal activities from endophytic actinomycetes

กิตติชัย เชื้อวชาญ¹ หทัยรัตน์ ราชนิยม¹ วรรัตน์ เครือสุวรรณ¹ ณัฐริกา แสงกฤช² และ อรินทิพย์ ธรรมชัยพิเนต^{1*}

Kittichai Cheawchan¹, Hathairat Rachniyom¹, Worarat Kruasuwan¹, Nattika Saengkrit² and Arinthip Thamchaipenet^{1*}

¹หน่วยวิจัยแอคติโนมัยซีท ภาควิชาพันธุศาสตร์ คณะวิทยาศาสตร์ มหาวิทยาลัยเกษตรศาสตร์ กรุงเทพฯ 10900;

²ศูนย์นาโนเทคโนโลยีแห่งชาติ สำนักงานพัฒนาวิทยาศาสตร์และเทคโนโลยีแห่งชาติ ปทุมธานี 12120

¹Actinobacteria Research Unit, Department of Genetics, Faculty of Science, Kasetsart University, Bangkok 10900; ²National Nanotechnology Center (NANOTEC), National Science and Technology Development Agency (NSTDA), Pathumthani 12120

*Corresponding author: arinthip.t@ku.ac.th

บทคัดย่อ

แอคติโนมัยซีทเอนโดไฟต์ 99 สายพันธุ์ ที่แยกได้จากอ้อย สามารถย่อยไคติน (76%) และเส้นใยราได้ (61%) รวมทั้งสามารถยับยั้งการเจริญของรา *Fusarium moniliforme* (71%) และ *Colletotrichum falcatum* (83%) ที่ก่อโรคเหี่ยวเน่าแดงในอ้อย โดยพบสายพันธุ์ที่สามารถยับยั้งราได้มากกว่า 70% และมีหลายสายพันธุ์ที่สามารถยับยั้งการเจริญของราได้ทั้งสองชนิด (71%) เมื่อนำสายพันธุ์ที่สามารถย่อยไคตินในระดับสูง (วงใส ขนาด 0.7–1.2 ซม) 4 สายพันธุ์ มาตรวจหายีน *chi18A* และ *chi19* ด้วยวิธีพีซีอาร์ พบว่าทุกสายพันธุ์มียีนทั้งสองชนิด

ABSTRACT

Ninety-nine strains of endophytic actinomycetes isolated from sugarcane were investigated and found that they could hydrolyze chitin (76%), degrade fungal mycelium (61%) and inhibit *Fusarium moniliforme* (71%) and *Colletotrichum falcatum* (83%), the causative agents of red rot wilt disease in sugarcane. Some strains showed more than 70% inhibition and several strains could inhibit both fungi (71%). When 4 high activity chitinase producing strains (clear zone from 0.7–1.2 cm) were screened for *chi18A* and *chi19* genes by PCR, all of them carried both genes.

คำสำคัญ: ไคติเนส, ยับยั้งรา, แอคติโนมัยซีท, เอนโดไฟต์, อ้อย

Keywords: chitinase, antifungal, actinomycete, endophyte, sugarcane

INTRODUCTION

Endophytes are microorganisms that reside in plants and help plants to enhance growth and protect them from phytopathogens. Endophytic actinomycetes have been reported as promising biocontrol agents because of their property of degradation of fungal cell wall by chitinase (El-Tarabily and Sivasithamparan, 2006). Chitinase is the enzyme that breaks down glycosidic bonds of chitin which is the major component of fungal cell wall. Chitinases are classified into family 18 and 19 of glycosyl hydrolases (Henrissat and Bairoch, 1993). Chitinolytic abilities from actinomycetes have been revealed to play a major role for control plant pathogenic fungi (Joo, 2005).

In this study, chitinase activity was screened from endophytic actinomycetes isolated from sugarcane. Antifungal activity against *Fusarium moniliforme* and *Colletotrichum falcatum*, the causative agents of red rot wilt disease in sugarcane (Jaiswal *et al.*, 1998), was also determined. Genes involved in chitinases were also identified.

MATERIALS AND METHODS

1. Investigation of chitinase activity

Endophytic actinomycetes isolated from sugarcane were obtained from GKU culture collection. They were tested for chitinase activity on colloidal chitin agar (CAA; Nawani *et al.*, 2002) and fungal mycelium fragmented agar (Valois *et al.*, 1996) using *Fusarium moniliforme*. Radiance of clear zone was observed after incubation at 28 °C for 14 days.

2. Antifungal activity of endophytic actinomycetes against pathogenic fungi

Endophytic actinomycetes were cultured on starch casein agar (SCA). *F. moniliforme* and *Colletotrichum falcatum* were inoculated 6 cm away from each of the actinomycete colony. Inhibition zone was observed after incubation at 28 °C for 7 days. Percentage of inhibition was calculated.

3. Identification of chitinase genes

Genomic DNAs of chitinase producing endophytic actinomycetes were prepared according to the protocol described by Kieser *et al.* (2000). Chitinase genes were amplified by PCR using specific primers for *chi18A* (ATT107F, 5'-GACACCTGGGACCAGCCGCTG-3' and ATT107R, 5'-TAGAAGCCGAYGCCGAKSAGCA-3') and *chi19* (ATT092F, 5'-CAGTTCRACCARATGTTCCCG-3') and ATT092R, 5'-CGTTGATSGASC GGATSGTCT-3').

RESULTS AND DISCUSSION

1. Chitinase activity from endophytic actinomycetes

Seventy-five strains from 99 of endophytic actinomycetes showed chitinase activity with various radiance of clear zone from 0.1–1.2 cm on colloidal chitin agar (76%) (Fig. 1a-c) after 14 days of incubation. Amongst these, 11 strains showed high activity level (0.7–1.2 cm). When the strains were tested on mycelium fragmented agar, 61 strains showed clear zone at average radiance about 0.1–1.3 cm (60%) (Fig. 1h-i) with 15 strains high activity level (0.7–1.3 cm).

2. Antifungal activity of endophytic actinomycetes

Seventy strains of 99 endophytic actinomycetes showed antifungal activity against *F. moniliforme* above 5% inhibition (71%) and 6 of the positive strains showed highest activity (more than 70% inhibition). Whereas, 82 strains showed activity against *C. falcatum* above 5% inhibition (83%) and 6 of the positives showed highest activity (more than 70% inhibition). In addition, 70 strains of the endophytes carried antifungal activity against both fungi (71%) (Fig. 1d-g).




Figure 1 Chitinase activity of endophytic actinomycetes on colloidal chitin agar (a-c). Zones of fungal inhibition against *F. moniliforme* (d), *F. moniliforme* negative control (e), against *C. falcatum* (f) *C. falcatum* negative control (g). Fungal mycelium degradation activity on *F. moniliforme* mycelium fragmented agar (h-i).

3. Chitinase producing endophytic actinomycetes

Endophytic actinomycetes in this study could be divided into three groups. The first group contains strains that carried high chitinase activity. This group could hydrolyzed chitin and degraded fungal mycelium but showed low antifungal activity. The second group contains low chitinase activity but high antifungal activity. The antagonism of this group is likely to be involved with the production of antifungal compounds. The last group contains high chitinase activity and high antifungal activity. The mechanism of this group may implicate in both of the previous mechanisms.

4. Identification of chitinase genes

Genomic DNAs of selected high chitinase producing strains from the first group proposed above namely, *Streptomyces* sp. GKU816, GKU853, GKU893 and a non-streptomycete strain GKU887 were amplified using specific primers for *chi18A* and *chi19* genes. The expected PCR products of 587 bp from *chi18A* and 483 bp for *chi19* were obtained from each strain (Fig. 2). The results were in agreement with previous report that chitinase gene family 18 and 19 were found in *Streptomyces coelicolor* (Kawase et al., 2006).




Figure 2 PCR products of *chi18A* and *chi19* genes. Lane M, DNA ladder; Lane 1-4, *chi18A* of GKU816, GKU853, GKU887 and GKU893; lane 5-8, *chi19* of GKU816, GKU853, GKU887 and GKU893.

CONCLUSION

Endophytic actinomycetes isolated from sugarcane revealed a good source for chitinase and antifungal activities against *F. moniliforme* and *C. falcatum*, the causative agents of red rot wilt disease in sugarcane. Further study on application of those strains as biocontrol agents in sugarcane is on the way.

ACKNOWLEDGEMENTS

This research was supported by TRF-Master research grant and Thailand Research fund.

REFERENCES

- El-Tarabily KA, Sivasithamparam K. Non-streptomycete actinomycetes as biocontrol agents of soil-borne fungal plant pathogens and as plant growth promoters. *Soil Biol Biochem.* 2006;38:1505-1520.
- Henrissat B, Bairoch A. New families in the classification of glycosyl hydrolases based on amino acid sequence similarities. *J Biochem.* 1993;293:781-788.
- Jaiswal AK, Rao GP, Pandey OP, Sengupta SK. Efficacy of organophosphorus derivatives against fungal pathogens of sugarcane. *J Agric Food Chem.* 1998;46:1609-1613.
- Joo GJ. Purification and characterization of an extracellular chitinase from the antifungal biocontrol agent *Streptomyces halstedii*. *Biotechnol Lett.* 2005;27:1483-1486.
- Kawase T, Yokokawa S, Saito A, Fujii T, Nikaidou N, Miyashita K, Watanabe T. Comparison of enzymatic and antifungal properties between family 18 and 19 chitinases from *S. coelicolor* A3(2). *Biosci Biotechnol Biochem.* 2006;70:988-998.
- Kieser T, Bibb MJ, Buttner MJ, Chater KF, Hopwood DA. *Practical Streptomyces Genetics.* 2000. The John Innes Foundation, Norwich.
- Nawani NN, Kapadnis BP, Das AD, Rao AS, Mahajan SK. Purification and characterization of a thermophilic and acidophilic chitinase from *Microbispora* sp. V2. *J Appl Microbiol.* 2002;93:965-975.
- Valois D, Fayad F, Barasubuye T, Garon M, Brezinski C, De'ryR, Beaulieu C. Glucanolytic actinomycetes antagonistic to *Phytophthora fragariae* var. *rubi*, the causal agent of raspberry root rot. *Appl Environ Microbiol.* 1996;62:1630-1635.