

## Occurrence of the Invasive Crab Species *Callinectes sapidus* Rathbun, 1896, in NW Greece

**Costas PERDIKARIS<sup>1</sup>, Evangelos KONSTANTINIDIS<sup>1</sup>, Evangelia GOUVA<sup>2</sup>,  
Anna ERGOLAVOU<sup>3</sup>, Dimitris KLAUDATOS<sup>4</sup>, Cosmas NATHANAILIDES<sup>5,6,\*</sup>  
and Ioannis PASCHOS<sup>2</sup>**

<sup>1</sup>Department of Fisheries, Regional Unit of Thesprotia, Region of Epirus, P. Tsaldari 18, 46100 Igoumenitsa, Greece

<sup>2</sup>Department of Agricultural Technology, Technological Educational Institute of Epirus, Kostakioi, 47100 Arta, Greece

<sup>3</sup>Karamanli 7, 46200 Paramythia, Greece

<sup>4</sup>Hellenic Centre of Marine Research, Athens, Helliniko, Greece

<sup>5</sup>Department of Fisheries and Aquaculture Technology, Technological Educational Institute of Western Greece, 30200 Messolonghi, Greece

<sup>6</sup>Laboratory of Ichthyology and Ichthyopathology, School of Veterinary Medicine, University of Thessaly, 43100 Karditsa, Greece

(\*Corresponding author's e-mail: cosmasfax@yahoo.com)

Received: 8 January 2015, Revised: 7 April 2015, Accepted: 6 May 2015

### Abstract

The present work aims to review and describe the current status of the invasive species *Callinectes sapidus* Rathbun, 1896, along the Ionian coastal zone of Greece and to assess its invasive potential. Blue crab has a long invasion history in the Mediterranean Sea, but the available data on the species occurrence in the Ionian sub-region are scarce and fragmented. The proximity of most of the estuaries and lagoons to local ports and to the shipping routes, as well as the swimming/dispersal ability of the adults of the species, indicate that range expansion will likely continue. The invasiveness risk of the species in the region was estimated with the use of a decision support tool (Marine Invertebrate Invasiveness Screening Kit, version 1.19). The observed impacts are discussed, along with urgent mitigation priorities. Measures to limit the expansion of this invasive species may include the effective management of ballast waters and the targeted increase of fishing pressure on *C. sapidus* populations. The importance of ballast water management is further highlighted by the existence of numerous ports with the capacity to serve ships with ballast tanks in close proximity to the recipient ecosystems. Moreover, the Ionian Sea, which connects the Adriatic Sea to the rest of the Mediterranean, is a significant shipping route for the local, regional, and international seaborne trade.

**Keywords:** Invasive species, ecological threats, marine pollution

### Introduction

The blue crab, *Callinectes sapidus* Rathbun, 1896 (Brachyura: Portunidae), native to the western Atlantic coast, has invaded, at least since the beginning of the 20<sup>th</sup> century, European waters in multiple independent introductions via ballast water. Invasion history and success on a Pan-European level was recently reviewed for the Atlantic Ocean coasts, North, Baltic, Mediterranean, Black, and Azov Seas by Nehring [1]. Concerning the Mediterranean Sea, the species has been recorded as having positively established populations in 12 countries. This exceptionally high establishment rate in this region, as well

as the reported negative impacts towards the coastal invaded ecosystems and the related fishing activities, justified its ranking among the 100 worst invasive species in the Mediterranean [2].

Early records (the earliest dating back to 1948) from Greece refer exclusively to the Aegean Sea [3-5], particularly to the northern coasts where the species is currently abundant [6,7]. Contrary to this, until recently, limited data were available in the scientific literature on the presence of the species on the Ionian coasts, although its occurrence has been documented since 1949 in Italy [8-12], 2004 in Croatia [13,14], and 2008 in Albania, near the border with Montenegro [15]. Despite the low number of alien marine species on the Ionian coasts compared to the Aegean Sea [16], the former sub-region has a very rich variety of estuaries, lagoons, and transitional waters, providing open ecological niches and favorable climatic conditions (i.e. the lowest, highest, and annual mean water temperatures during 2013 on the north-western Ionian coast of Greece were 13.8, 27.1, and 19.6 °C, respectively; Perdikaris, unpub. data). These conditions match the biological requirements of *C. sapidus*, as the reproductive cycle is related to access to brackish water, and the first developmental stages need temperatures of above 20 °C [17].

Within the above context, the initial step of any management intervention has to rely on the earliest possible detection of the introduced species. This should be followed by the determination of the risk level it poses to the recipient water bodies, including socio-economic considerations [18]. Therefore, given the scarcity of zoogeographical data on *C. sapidus* on the Ionian coasts of Greece, the aim of the present work was to: a) review and depict the occurrence of the species in this particular sub-region; b) assess its invasiveness risk, and c) discuss any observed negative impacts on biodiversity and fisheries, in order to suggest adequate mitigation measures.

## Materials and methods

During 2012 - 2013, data were retrieved from peer reviewed published papers in the scientific literature, private communications and direct contacts with ichthyologists (in Regional Fisheries Departments), and fishermen and members of fishermen's associations and estuary/lagoon co-operatives along the entire length of the Ionian coasts (including the Islands of Corfu and Lefkas). Data on species occurrence were subsequently inserted into a GIS software (ArcMap 10.1, Esri, USA) to generate a distribution map, including all major and minor ports in the sub-region which have the minimum infrastructure to harbor ships with ballast tanks (i.e. above 300 gross registered tonnage - GRT). Finally, the invasive potential of the species was preliminarily assessed by one of us (CP) using the Marine Invertebrate Invasiveness Screening Kit (MI-ISK v1.19) (available at <http://www.cefas.defra.gov.uk/our-science/ecosystems-and-biodiversity/non-native-species/decision-support-tools.aspx>). The final score was based on the responses to 49 questions, related to topics such as domestication/cultivation, climate and distribution, invasiveness in a non-native range, undesirable traits, feeding guild, reproduction, dispersal mechanisms, and persistence attributes. Based on the responses, a numerical score was calculated according to the following thresholds: accept (score < 0 = low risk), evaluate (score 1 - 6 = medium risk), or reject (> 6 = high risk). The scoring was provisional, given that the tool is not calibrated yet.

## Results and discussion

*Callinectes sapidus* occurs in 9 regional units (former prefectures) along the Ionian coasts of Greece (including the islands of Corfu and Lefkas) and in the majority of the estuaries/lagoons of the sub-region (**Figure 1, Table 1**). The earliest records date back to the late 1990s and, in most cases, the species was detected during the 2000s.


The following key biological features and ecological traits of *C. sapidus* are exceptionally suitable for further range expansion in the Ionian sub-region. The enormous reproductive potential, e.g. the number of extruded eggs that has recently been calculated at  $1.96 \times 10^6$  [19], the early maturity, and fast growth reported for the northern Aegean Sea [6,7] suggests that the species has been successfully established in Greece, and efficiently exploits the available resources in brackish waters and transitional and coastal marine environments. This adaptation and the on-going invasion process is further assisted by its ability (at least in the native range) to move from shallow vegetated estuaries (juveniles) to deep and

non-vegetated estuaries (adults) during the development and the recruitment process at a spatial scale of  $10^4$  to  $10^5$  m [20]. The above swimming ability suggests that further invasion of new areas from the initial establishment sites should be expected not only through passive dispersal (i.e. transportation of larvae in ballast waters), but also by energetic swimming ( $> 1$  m sec $^{-1}$ ) [21]. Subsequently, when established, the effective competition for space and resources, aided by the large size, the armored exoskeleton, its hardness, and its polytrophic feeding habits [17], are crucial attributes for dominance in the trophic webs. Finally, large juveniles (i.e.  $> 75$  mm) and adults are vulnerable to predators (piscivorous fish and birds), mainly during moulting (i.e. soft-shell stage) [22].

The application of the MI-ISK risk identification tool, although still non-calibrated, suggested a high risk of invasiveness (i.e. total score of 28), as the species was rejected for introduction/translocation in the risk assessment area. Concerning score partitioning, the scores for biogeography, undesirable attributes, and biology/ecology were 10, 9, and 9, respectively. This is the first attempt to assess the invasiveness potential of *C. sapidus*, following the assessment of freshwater crayfish species in Italy and Greece [23,24] and of Chinese mitten crab *Eriocheir sinensis* Milne-Edwards, 1853, in the Serbian section of the River Danube [25].

**Table 1** Occurrence, abundance and estimated date of first presence of blue crab *Callinectes sapidus* Rathbun, 1896, in the Ionian coast in Greece.

| Regional unit  | Ecosystem  | Estimated date of first presence | Observed or likely impacts (negative/positive)   | Source   |
|----------------|--|----------------------------------|--|--|
| Corfu          | Antinioti lagoon   | Late 2000s                       | Small population commercially exploited  | Fisheries Department in Corfu; [31]                                  |
|                | Chalkiopoulos lagoon   | Late 2000s                       | Unknown  | Fisheries Department in Corfu  |
| Thesprotia     | Voda estuary/mouth of River Kalamas  | Late 1990s                       | Consuming netted fish and cutting gill nets  | [19,37]  |
|                | Richo/Vatatsa lagoon complex   | Late 2000s                       | Unknown  | Lagoon co-operative  |
|                | Bay of Igoumenitsa   | Late 2000s                       | Consuming netted fish and cutting gill nets  | Fishermen; Local Fishermen Association                               |
| Preveza/Arta   | Lagoons complex of Rodia, Tsoukalio, and Logarou in the northern part of the Amvrakikos Gulf | Early 2000s                      | Unknown  | Fisheries Department in Preveza; Amvrakikos Wetlands Management Body |
| Lefkas         | Lefkas lagoon and in nearby coastal zone   | 2000s                            | Unknown  | Fishermen; Fishmongers   |
| Etoloakarnania | Tholi lagoon and sporadically in most lagoons and estuaries of the Mesolongi complex         | Early 2000s                      | Unknown  | Fisheries Department in Mesolongi                                    |
|                | Mpouka lagoon/mouth of River Achelous  | Early 2000s                      | Unknown  | Fisheries Department in Mesolongi                                    |
| Achaia<br>Ilia | Pappas lagoon  | Unknown                          | Unknown  | Fisheries Department in Patra  |
|                | Kotychi lagoon   | Late 1990s                       | Consuming fish and cutting gill nets; harvesting difficulties of fish in the fixed structures of the lagoons due to the presence of high number of crabs | Fishermen; Lagoon co-operative                                       |
| Messinia       | Mouth of River Pamisos   | 2010s?                           | Unknown  | [38]   |



**Figure 1** Map of occurrence of *Callinectes sapidus* on the Ionian coasts of Greece. Black and open circles with dots indicate major and minor ports, respectively. Invaded coastal ecosystems are shaded in black. 1: Antinioti lagoon; 2: Chalkiopoulos lagoon; 3: Voda estuary/mouth of River Kalamas; 4: Vatatsa lagoon; 5: Richo lagoon; 6: Bay of Igoumenitsa; 7: Logarou lagoon; 8: Tsoukalio lagoon; 9: Rodia lagoon; 10: Lefkas lagoon and nearby coast; 11: Mpouka lagoon/mouth of River Acheloos; 12: Tholi lagoon and lagoons and estuaries of the Mesolongi complex; 13: Pappas lagoon; 14: Kotychi lagoon; 15: Mouth of River Pamisos.

Considering the scarcity of documented impacts in the Mediterranean region, the species is likely in competition with other crabs [9]. This invasive species could compete with other crab species, increasing, for example, the natural mortality, distribution, and dynamics of indigenous crabs [26]. In fact, a species may be important for the economy of a region in which it is indigenous, but its expansion-invasion in other ecosystems-regions may result in considerable economic and ecological damages [27,28]. Observed impacts are related to severe damage in fishing nets when *C. sapidus* try to feed on fish caught in them [29]. The species has also been suggested as a possible bio-accumulator of toxins and a carrier of strains of the human cholera [1]. In this work, impacts were reported in three ecosystems (**Table 1**). In these ecosystems, blue crabs cause problems to fishing by tearing nets, and to fish harvesting when the crabs aggregate in the fixed harvesting installations in the lagoon mouth. However, there is no solid evidence to support negative impacts either on biodiversity (e.g. direct predation, displacement, or growth suppression of other invertebrates, such as native crabs or sea cucumber populations), or on local mussel farms in the Gulf of Amvrakikos in West Greece.

Nevertheless, the trade of the species may result in some economic benefits. Currently, the exploitation of the species is very limited and localized, compared, for instance, to northern Greece (annual landings of 50 - 80 tons during 2010 - 2011 in the auction market of Thessaloniki; [7]), or to the trap fisheries of Turkey [30]. The species is used in the local cooking of Corfu [31] where it is sold mainly in street markets at retail prices of €3 - 5 per kg and, although it is sporadically seen in fishmonger stores in many cities (e.g. in Patra, Igoumenitsa, Arta, Lefkada), it is rather unknown to the consumers in the continental part of the Ionian sub-region. This could be due to the recent invasion history of the species and the lack of a crab-eating tradition.

If a marine invasive species is not detected shortly after arrival, and becomes widely distributed, there are no proven techniques to eradicate it [32]. This is exactly the case for *C. sapidus* in Greece, and further range expansion in all suitable transitional ecosystems in the Ionian Sea is likely inevitable, given that established populations are already numerous and widespread. However, mitigation (i.e. control of the species so that the provision of essential ecosystem services is continued in its presence) would be a sound option. For *C. sapidus*, which has few year classes spawning, the effects of overfishing could have a serious effect on population recruitment [33]. Constant pressure on these populations by crabbing (similarly to certain overexploited US populations [34]) could be an effective method of maintaining local abundance at as low a level as possible. Moreover, the possibility of employing *C. sapidus* as a fish feed additive, to improve color enhancement, and as flavor attractants, should be experimentally assessed. For example, natural astaxanthin production utilizes by-products of Antarctic krill, crab, crayfish and shrimp meals, and the content of carotenoids in crab meals from selected sources may reach 1300 mg/kg [35].

The importance of ballast water management and the use of filters in ships is highlighted by a) the existence of numerous (23) ports with the capacity to serve ships with ballast tanks in close proximity to the recipient ecosystems (Figure 1), and b) the significance of the Ionian shipping route (i.e. in terms of increased traffic volume) to the local, regional and international seaborne trade, connecting the Adriatic Sea to the rest of the Mediterranean. Accordingly, the implementation of the Ballast Water Management Convention (BWMC) [36] by all countries in the Adriatic-Ionian region is an urgent priority, in order to control the passive pathway of new introductions [37,38] (e.g. all ships should implement a ballast water management plan, carry a ballast water record book, and carry out ballast water management procedures to a given standard). Within the above context, the installation of filters could be a practical and cost-effective option compared to other methods, such as heat treatment, chemical treatment, and ultraviolet light, which are expensive and with certain drawbacks. Currently, the most common treatment method applied is ballast water exchange at sea, but the efficacy of this method is unknown. The volume of a ballast tank is pumped 3 times through the tank. By this, it is expected to lead to 95 % exchange of the original ballast water, replacing it with oceanic ballast that poses little threat to coastal ecosystems [32].

## Conclusions

The data of the present work suggest that *C. sapidus* has invaded a significant amount of estuaries, lagoons, and transitional waters along the entire coastal zone of the Ionian Sea in Greece. This range expansion will likely continue due to active (swimming) spread and passive (ballast water) transportation. Concerning the negative effects, these are mainly related to damages in fishing nets and fish harvesting difficulties in lagoons. Given its high invasiveness risk, mitigation measures may include the increase of fishing pressure targeting the species, the effective management of ballast waters, and the utilization of the species as an additive for animal feeds.

## Acknowledgements

The authors would like to thank all the participants in the survey and, particularly, S. Ginis (State Fisheries Department in Corfu), C. Akovitiotis (State Fisheries Department in Preveza), S. Konstas (Amvrakikos Wetlands Management Body), V. Dimitriou (State Fisheries Department in Mesolongi), I. Ravrasopoulos (State Fisheries Department, Patra), and the fishermen D. Papadimitriou and P. Karkavitsas, for assistance in collecting the data from the fishermen in the region.

## References

- [1] S Nehring. *Invasion History and Success of the American Blue Crab Callinectes sapidus in European and Adjacent Waters*. In: BS Galil, PF Clark and JT Carlton (eds.). *In the Wrong Place-Alien Marine Crustaceans: Distribution, Biology and Impacts. Invading Nature - Springer Series in Invasion Ecology*, 2011, p. 607-24.
- [2] N Streftaris and A Zenetos. Alien marine species in the Mediterranean - the 100 “Worst Invasives” and their impact. *Mediterr. Mar. Sci.* 2006; **7**, 87-118.
- [3] C Serbetis. Un nouveau crustacé commestible en mer Egee *Callinectes sapidus* Rath. (Decapod brach.). *Proc. Gen. Fish. Counc. Medit.* 1959; **5**, 505-7.
- [4] R Kinzelbach. Die Blaue Schwimmkrabbe (*Callinectes sapidus*), ein Neubürger im Mittelmeer. *Nat. Mus.* 1965; **95**, 293-6.
- [5] LB Holthuis. Enkele interessante Nederlandse crustacean. Bijdragen tot de faunistiek van Nederland. *I. Zool. Bijdr. Leiden* 1969; **11**, 34-48.
- [6] K Kevrekidis, K Avramoglou, J Efsthathiadis and C Chintiroglou. Relative growth of the blue crab *Callinectes sapidus* in Thermaikos Gulf (Methoni Bay): Preliminary results. In: Proceedings of the 15<sup>th</sup> Pan-Hellenic Congress of Ichthyologists, Thessaloniki, Greece, 2013, p. 109-12.
- [7] K Kevrekidis, C Antoniadou, K Avramoglou, J Efsthathiadis and C Chintiroglou. Population structure of the blue crab *Callinectes sapidus* in Thermaikos Gulf (Methoni Bay). In: Proceedings of the 15<sup>th</sup> Pan-Hellenic Congress of Ichthyologists, Thessaloniki, Greece, 2013, p. 113-6.
- [8] L Mizzan. Presence of swimming crabs of the genus *Callinectes* (Stimpson) (Decapoda, Portunidae) in the Venice Lagoon (North Adriatic Sea - Italy): First record of *Callinectes danae* Smith in European waters. *Boll. Mus. Civ. Stor. Nat. Venezia* 1993; **42**, 31-43.
- [9] R Gennaio, G Scordella and M Pastore. Occurrence of blue crab *Callinectes sapidus* (Rathbun, 1986 Crustacea, Brachyura), in the Ugento ponds area (Lecce, Italy). *Thalas. Salent.* 2006; **29**, 29-39.
- [10] M Florio, P Breber, T Scirocco, A Specchiulli, L Cilenti and L Lumare. Exotic species in Lesina and Varano lakes: Gargano National Park (Italy). *Transit. Waters Bull.* 2008; **2**, 69-79.
- [11] L Castriota, F Andaloro, R Costantini and AD Ascentiis. First record of the Atlantic crab *Callinectes sapidus* Rathbun, 1896 (Crustacea: Brachyura: Portunidae) in Abruzzi waters, central Adriatic Sea. *Acta Adriat.* 2012; **53**, 467-71.
- [12] G Mancinelli, L Carrozzo, ML Costantini, L Rossi, G Marini and M Pinna. Occurrence of the Atlantic blue crab *Callinectes sapidus* Rathbun, 1896 in two Mediterranean coastal habitats: Temporary visitor or permanent resident? *Estuar. Coast. Shelf S.* 2013; **135**, 46-56.

- [13] V Onofri, J Dulčić, A Conides, S Matić-Skoko and B Glamuzina. The occurrence of the blue crab, *Callinectes sapidus* Rathbun, 1896 (Decapoda, Brachyura, Portunidae) in the eastern Adriatic (Croatian coast). *Crustaceana* 2008; **81**, 403-9.
- [14] J Dulčić, P Tutman, S Matić-Skoko and B Glamuzina. Six years from first record to population establishment: the case of the blue crab, *Callinectes sapidus* Rathbun, 1896 (Brachyura, Portunidae) in the Neretva River delta (south-eastern Adriatic Sea, Croatia). *Crustaceana* 2011; **84**, 1211-20.
- [15] S Beqiraj and L Kashta. The establishment of blue crab *Callinectes sapidus* Rathbun, 1896 in the Lagoon of Patok, Albania (south-east Adriatic Sea). *Aquat. Invas.* 2010; **5**, 219-21.
- [16] MA Pancucci-Papadopoulou, A Zenetos, M Corsini-Foka and CY Politou. Update of marine alien species in Hellenic waters. *Mediterr. Mar. Sci.* 2005; **6**, 147-58.
- [17] MR Millikin and AB Williams. *Synopsis of Biological Data on the Blue Crab Callinectes sapidus Rathbun*. FAO Fisheries Synopsis, University of California Libraries, 1984, p. 1-39.
- [18] RE Gozlan, JR Britton, I Cowx and GH Copp. Current knowledge on non-native freshwater fish introductions. *J. Fish Biol.* 2010; **76**, 751-86.
- [19] M Thessalou-Legaki, O Aydogan, P Bekas, G Bilge, YO Boyaci, E Brunelli, V Circosta, F Crocetta, F Durucan, M Erdem, A Ergolavou, H Filiz, F Fois, E Gouva, K Kapiris, S Katsanevakis, Z Kljajic, E Konstantinidis, G Konstantinou, D Koutsogiannopoulos, S Lamon, V Macic, R Mazzette, D Meloni, A Mureddu, I Paschos, C Perdikaris, F Piras, D Poursanidis, AA Ramos-Espala, A Rosso, P Sordino, E Sperone, A Sterioti, E Taskin, F Toscano, S Tripepi, L Tsiakkios and A Zenetos. New mediterranean biodiversity records (December 2012). *Mediterr. Mar. Sci.* 2012; **13**, 312-27.
- [20] BM Gillanders, KW Able, JA Brown, DB Eggleston and PF Sheridan. Evidence of connectivity between juvenile and adult habitats for mobile marine faunas: An important component of nurseries. *Mar. Ecol. Prog. Ser.* 2003; **247**, 281-95.
- [21] CP Spirito. An analysis of swimming behavior in the portunid crab *Callinectes sapidus*. *Mar. Behav. Physiol.* 1972; **1**, 261-76.
- [22] AH Hines, AM Haddon and LA Wiechert. Guild structure and foraging impact of blue crabs and epibenthic fish in a subestuary of Chesapeake Bay. *Mar. Ecol. Prog. Ser.* 1990; **67**, 105-26.
- [23] I Papavlasopoulou, C Perdikaris, L Vardakas and I Paschos. Enemy at the gates: Introduction potential of non-indigenous freshwater crayfish in Greece via the aquarium trade. *Cent. Eur. J. Biol.* 2014; **9**, 11-8.
- [24] E Tricarico, LVilizzi, F Gherardi and GH Copp. Calibration of FI-ISK, an invasiveness screening tool for nonnative freshwater invertebrates. *Risk Anal.* 2010; **30**: 285-92.
- [25] D Škraba, A Tošić, D Miličić, V Nikolić and P Simonović. Invasiveness assessment of the Chinese mitten crab *Eriocheir sinensis* (H. Milne Edwards, 1853) in the Serbian section of the river Danube. *Arch. Biol. Sci.* 2013; **65**, 353-8.
- [26] A Sawusdee and A Songrak. Population dynamics and stock assessment of blue swimming crab (*Portunus pelagicus* Linnaeus, 1758) in the coastal area of Trang Province, Thailand. *Walailak J. Sci. & Tech.* 2011; **6**, 189-202.
- [27] HJ Li, CB He, Q Yang, ZG Shan and XG Gao. Characterization of single nucleotide polymorphisms from expressed sequence tags of chinese mitten crab *Eriocheir sinensis*. *Aquat. Biol.* 2010; **11**, 193-9.
- [28] D Klaoudatos and K Kapiris. *Alien Crabs in the Mediterranean Sea*. In: C Ardoxin (ed.). Crabs: Global Diversity, Behaviour and Environmental Threats. Nova Publishers, USA, 2014, p. 101-60.
- [29] A Zenetos, MA Pancucci-Papadopoulou, S Zogaris, E Papastergiadou, L Vardakas, K Aligizaki and A Economou. Aquatic alien species in Greece (2009): Tracking sources, patterns and effects on the ecosystem. *J. Biol. Res. Thessalon.* 2009; **12**, 135-72.
- [30] H Atar, M Ölmez, S Bekcan and S Seçer. Comparison of three different traps for catching blue crab (*Callinectes sapidus* Rathbun 1896) in Beymelek Lagoon. *Turk. J. Vet. Anim. Sci.* 2009; **26**, 1145-50.
- [31] M Bilecenoglu, JEF Alfaya, E Azzurro, R Baldacconi, YO Boyaci, V Circosta, LJV Compagno, F Coppola, A Deidun, H Durgham, F Durucan, D Ergüden, FA Fernández-Álvarez, P Gianguzzza, G Giglio, M Gökoğlu, M Gürlek, S Ikhtiyar, H Kabasakal, PK Karachle, S Katsanevakis, D

- Koutsogiannopoulos, E Lanfranco, P Micarelli, Y Özvarol, L Pena-Rivas, D Poursanidis, J Saliba, E Sperone, D Tibullo, F Tiralongo, S Triepi, C Turan, P Vella, MB Yokeş and B Zava. New Mediterranean Marine biodiversity records (December, 2013). *Mediterr. Mar. Sci.* 2013; **14**, 463-80.
- [32] N Bax, A Williamson, M Aguero, E Gonzalez and W Geeves. Marine invasive alien species: A threat to global biodiversity. *Mar. Policy* 2003; **27**, 313-23.
- [33] CM Jones, JR McConaugha, PJ Geer and MH Prager. Estimates of spawning stock size of blue crab, *Callinectes sapidus*, in Chesapeake Bay, 1986-1987. *Bull. Mar. Sci.* 1990; **46**, 159-69.
- [34] AF Sharov, JH Vølstad, GR Davis, BK Davis, RN Lipcius and MM Montane. Abundance and exploitation rate of the blue crab (*Callinectes sapidus*) in Chesapeake Bay. *Bull. Mar. Sci.* 2003; **72**, 543-65.
- [35] SK Gupta, AK Jha, AK Pal and G Venkateshwarlu. Use of natural carotenoids for pigmentation in fishes. *Nat. Prod. Rad.* 2007; **6**, 46-9.
- [36] International Convention or the Control and Management of Ships' Ballast Water and Sediments (BWM). Available at: <http://www.imo.org/About/Conventions/ListOfConventions/Pages/International-Convention-for-the-Control-and-Management-of-Ships%27-Ballast-Water-and-Sediments-%28BWM%29.aspx>, accessed November 2014.
- [37] C Perdikaris. The presence of blue crab (*Callinectes sapidus* Rathbun, 1896) in Greece. In: Proceedings of the 15<sup>th</sup> Pan-Hellenic Congress of Ichthyologists, Thessaloniki, Greece, 2013, p. 447.
- [38] A Eleftheriou, E Anagnostopoulou-Visilia, E Anastasopoulou, AS Ates, NEI Bachari, L Cavas, C Cevik, M Ulha, F Cevik, AL Delos, OB Derici, D Erguden, N Fragopoulou, A Giangrande, T Goksan, C Gravili, M Gurlek, A Hattour, K Kapiris, P Kouraklis, S Lamouti, E Prato, L Papa, G Papantonio, I Parlapiano, D Poursanidis, C Turan and D Yaglioglu. New mediterranean biodiversity records (December 2011). *Mediterr. Mar. Sci.* 2012; **12**, 491-508.